



## Prospects for Indirect Detection of Neutralino Dark Matter

Jonathan L. Feng<sup>a</sup>, Konstantin T. Matchev<sup>b</sup>, and Frank Wilczek<sup>a</sup>

<sup>a</sup> *School of Natural Sciences, Institute for Advanced Study  
Princeton, NJ 08540, U.S.A.*

<sup>b</sup> *Theoretical Physics Department, Fermi National Accelerator Laboratory  
Batavia, IL 60510, U.S.A.*

### Abstract

Dark matter candidates arising in models of particle physics incorporating weak scale supersymmetry may produce detectable signals through their annihilation into neutrinos, photons, or positrons. A large number of relevant experiments are planned or underway. The ‘logically possible’ parameter space is unwieldy. By working in the framework of minimal supergravity, we can survey the implications of the experiments for each other, as well as for direct searches, collider searches, low-energy experiments, and naturalness in a transparent fashion. We find that a wide variety of experiments provide interesting probes. Particularly promising signals arise in the mixed gaugino-Higgsino region. This region is favored by low-energy particle physics constraints and arises naturally from minimal supergravity due to the focus point mechanism. Indirect dark matter searches and traditional particle searches are highly complementary. In cosmologically preferred models, if there are charged superpartners with masses below 250 GeV, then some signature of supersymmetry must appear *before* the LHC begins operation.

*Submitted to Phys. Rev. D*

## I. INTRODUCTION

It is now well established that luminous matter makes up only a small fraction of the mass of the observed universe. The evidence for dark matter is both astrophysical and cosmological [1]. Such evidence requires only that dark matter is gravitationally interacting. However, additional constraints, especially the success of light-element cosmonucleosynthesis calculations, strongly disfavor the possibility that dark matter is composed solely of baryons [2], and so some form of matter foreign to our everyday world is required. The dark matter problem is therefore also an important problem for particle physics, as particle physics both suggests promising possibilities and imposes stringent constraints.

Neutralinos are well-motivated candidates to provide much or all of the non-baryonic dark matter. An effectively stable particle is a generic component of models with weak-scale supersymmetry. This particle is the lightest supersymmetric particle (LSP), and is typically the neutralino, a mixture of the superpartners of Higgs and electroweak gauge bosons. Particle physics considerations alone require the neutralino to be electrically neutral, effectively stable (assuming  $R$ -parity conservation, which is also motivated by the need to forbid too-rapid proton decay), and weakly interacting, with mass of order 100 GeV (required, as we shall quantify below, if supersymmetry naturally protects the electroweak scale from large radiative corrections). Remarkably, these properties are consistent with the possibility that the thermal relic density of neutralinos makes up most of the missing mass of the universe [3,4].

Unfortunately, these properties also guarantee that neutralinos are practically impossible to observe in collider experiments directly. They pass through collider detectors without interacting. Existing bounds on neutralinos therefore rely on model-dependent correlations between their properties and those of other supersymmetric particles. At present, in minimal supergravity, the LEP experiments constrain  $m_\chi \gtrsim 40$  GeV [5]. In the next several years, Run II of the Tevatron at Fermilab and, eventually, the Large Hadron Collider at CERN will provide more powerful collider probes.

If neutralinos make up a significant portion of the halo dark matter, many additional avenues for their detection open up. They may deposit energy as they scatter off nuclei in detectors. We have investigated the prospects for direct detection in a companion article [6], where we emphasized the importance and promise of a mixed gaugino-Higgsino regime, previously neglected. Here we will study the possibility of detecting neutralinos indirectly by looking for evidence of their annihilation [7]. In the next five years, an astounding array of experiments will be sensitive to the various potential neutralino annihilation products. These include under-ice and underwater neutrino telescopes (AMANDA, NESTOR, ANTARES), atmospheric Cherenkov telescopes (STACEE, CELESTE, ARGO-YBJ, MAGIC, HESS, CANGAROO, VERITAS), space-based  $\gamma$  ray detectors (AGILE, AMS/ $\gamma$ , GLAST), and anti-matter/anti-particle experiments (PAMELA, AMS). In many cases, these experiments will improve current sensitivities by several orders of magnitude.

In this paper we evaluate the prospects for neutralino dark matter discovery through indirect detection. The neutralino signals depend on many unknown parameters. At the same time, an abundance of theoretical and experimental information from particle physics can be brought to bear. The implications of traditional particle physics searches for dark matter searches, and *vice versa*, are already significant, and promise to become much stronger

over the next few years. One of our main conclusions is that in a class of particle physics models favored by current particle physics constraints, astrophysical signals are especially enhanced.

Previous discussions of indirect neutralino detection fall rather sharply into two schools. Several previous works are based on specific high energy models [8,9], including a number in the framework of minimal supergravity [10–14]. As we will recall below, minimal supergravity incorporates several desirable features, including the radiative breaking of electroweak symmetry and the possibility, suggested by gauge coupling unification [15], of perturbative extrapolation to large energy scales. Previous studies in minimal supergravity have concluded that only Bino-like dark matter is allowed by particle physics constraints. Such dark matter necessarily implies highly suppressed dark matter signals, as we will see. These studies, and their somber conclusions, have been criticized as products of overly restrictive particle physics assumptions [16]. Recently we have argued more specifically that, even in minimal models, these studies failed to examine a very well-motivated regime of parameters, and that for this reason their conclusions are overly pessimistic [6].

At the other extreme, several studies scan over a large set of weak-scale supersymmetry parameters and consider values for these parameters as large as 50 TeV. (See, for example, Refs. [16–20].) These studies, and others, bring a high level of sophistication to the evaluation of astrophysical effects on dark matter signals. In this regard, we will have nothing to add, but we will incorporate many of the most accurate recent results in our study.

From a particle physics perspective, this second group of studies is impressively general, but this generality is achieved at a cost. For example, extrapolating a given set of weak-scale parameters to higher scales, one may encounter such diseases as Landau poles or charge- or color-breaking minima. Models of this sort do not arise within a reasonable high energy framework. In addition, there is the practical difficulty that the proliferation of free parameters implies that results cannot be presented in a systematic, yet transparent, fashion. Typically they are displayed as scatter plots after scanning over all parameters. It is nearly impossible, from such plots, to determine the dependence of the signal rates on the underlying physical parameters. Dedicated correlation plots have been used to highlight a few of the relations between dark matter detection experiments, but even the most general implications of these experiments for collider searches, electric dipole moments, anomalous magnetic moments, proton decay, flavor violation, and other searches for supersymmetry are very hard to discern. Finally, it can be expected that supersymmetry parameters, such as the  $\mu$  and gaugino mass parameters, of order 50 TeV will require fine-tuning of the order of 1 part in  $10^6$  to produce the observed electroweak scale. (While it is impossible to speak of naturalness without first specifying a mechanism of electroweak symmetry breaking, at present all concrete models display this feature.) Such large fine-tuning destroys one of the main motivations for considering supersymmetric extensions of the standard model in the first place. Lacking both a systematic framework and a systematic presentation, it is impossible to see how the expectations narrow when some naturalness criterion is imposed.

It may be impossible to satisfy both schools simultaneously. However, recent results suggest an appealing compromise. As has been emphasized in Refs. [21–23], a Bino-like LSP is *not* a robust prediction of minimal supergravity. We find that both Bino-like and mixed gaugino-Higgsino dark matter is possible. So simply by considering all of minimal supergravity parameter space carefully, as we will do here, we remove the most egregious

form of model dependence. At the same time, by staying within the confines of minimal supergravity we will be able to present results in an organized and comprehensive manner, so that correlations with all other supersymmetric signals are easily determined. As the experiments discussed below report results, it will be ever more interesting to see what models are being excluded or favored. The framework discussed here makes this possible.

Inclusion of the new gaugino-Higgsino LSP models in minimal supergravity is not just a formality. The region with mixed gaugino-Higgsino LSPs is now known to be robust and natural, given an objective definition of naturalness [21–23]. It yields cosmologically interesting relic densities [6], and is even favored by low energy constraints such as proton decay and electric dipole moments [24]. As we will see, *all* indirect detection signals are enhanced in this region. This lends increased interest to indirect dark matter searches, since large — possibly spectacular — rates are predicted within an attractive and simple high energy framework.

In the following section, we review a few essential results concerning neutralino dark matter in minimal supergravity with an emphasis on the new gaugino-Higgsino LSP region. We then consider each of three promising signals in the following sections: upward-going muons from neutrinos in Sec. III, photons in Sec. IV, and positrons in Sec. V. In Sec. VI we compare these to direct dark matter and traditional particle physics searches, and in Sec. VII we summarize our results.

## II. NEUTRALINO DARK MATTER AND ITS ANNIHILATION

The lightest neutralino is the LSP in many supersymmetric models. Assuming  $R$ -parity conservation to prevent too-rapid proton decay through dimension-four operators, the LSP is effectively stable, and the neutralino is then an excellent candidate for cold dark matter.

The signals of neutralino dark matter are determined in large part by their composition. Neutralinos are mixtures of the superpartners of Higgs and electroweak gauge bosons. After electroweak symmetry breaking, these gauge eigenstates mix through the tree-level mass matrix

$$\begin{pmatrix} M_1 & 0 & -m_Z \cos \beta \sin \theta_W & m_Z \sin \beta \sin \theta_W \\ 0 & M_2 & m_Z \cos \beta \cos \theta_W & -m_Z \sin \beta \cos \theta_W \\ -m_Z \cos \beta \sin \theta_W & m_Z \cos \beta \cos \theta_W & 0 & -\mu \\ m_Z \sin \beta \sin \theta_W & -m_Z \sin \beta \cos \theta_W & -\mu & 0 \end{pmatrix} \quad (1)$$

in the basis  $(-i\tilde{B}, -i\tilde{W}^3, \tilde{H}_u^0, \tilde{H}_d^0)$ . The weak scale parameters entering this mass matrix are the Bino, Wino, and Higgsino mass parameters  $M_1$ ,  $M_2$ , and  $\mu$ , and the ratio of Higgs scalar vacuum expectation values  $\tan \beta = \langle H_u^0 \rangle / \langle H_d^0 \rangle$ . The lightest neutralino mass eigenstate is then determined by these parameters to be some mixture

$$\chi \equiv \chi_1^0 = a_1(-i\tilde{B}) + a_2(-i\tilde{W}^3) + a_3\tilde{H}_u^0 + a_4\tilde{H}_d^0. \quad (2)$$

We define the LSP gaugino fraction to be

$$R_\chi \equiv |a_1|^2 + |a_2|^2. \quad (3)$$

In the following, we refer to neutralinos with  $0.9 < R_\chi$  as gaugino-like,  $0.1 \leq R_\chi \leq 0.9$  as mixed gaugino-Higgsino, and  $R_\chi < 0.1$  as Higgsino-like.

The preceding discussion is model-independent, assuming only minimal field content. However, the minimal supersymmetric standard model is undoubtedly a low-energy effective theory of a more fundamental theory defined at some higher scale, such as the grand unified theory (GUT) or string scale. A simple realization of this idea is the framework of minimal supergravity, which is fully specified by the five parameters (four continuous, one binary)

$$m_0, M_{1/2}, A_0, \tan \beta, \text{sgn}(\mu) . \quad (4)$$

Here,  $m_0$ ,  $M_{1/2}$ , and  $A_0$  are the universal scalar mass, gaugino mass, and trilinear scalar coupling. They are assumed to arise through supersymmetry breaking in a hidden sector at the GUT scale  $M_{\text{GUT}} \simeq 2 \times 10^{16}$  GeV. The hidden-sector parameters then determine all the couplings and masses of the weak scale Lagrangian through renormalization group evolution. In particular, electroweak symmetry is broken radiatively by the effects of a large top quark Yukawa coupling, and the electroweak scale is determined in terms of supersymmetry parameters through

$$\frac{1}{2}m_Z^2 = \frac{m_{H_d}^2 - m_{H_u}^2 \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2 . \quad (5)$$

Equation (5) receives corrections beyond tree-level; in our work we include all one-loop effects in the Higgs potential [25]. We also use two-loop renormalization group equations [26] with one-loop threshold corrections [25,27] and calculate all superpartner masses to one-loop [25]. All of the qualitative features to be described below are present, however, also for one-loop renormalization group equations.

Minimal supergravity is, of course, by no means the most general allowed framework. It is worth noting, however, that the assumptions most relevant for dark matter, namely, the universality of gaugino and scalar masses, are motivated not only by their simplicity, but also by concrete experimental facts. The case for gaugino mass unification is especially powerful. In the minimal supersymmetric standard model, the three gauge couplings, when evolved to high scales, meet with high precision at a point, the GUT scale [15]. If the standard model is unified in a grand unified gauge theory, one typically obtains also gaugino mass unification. The unification of couplings calculation also distinguishes  $M_{\text{GUT}} \simeq 2 \times 10^{16}$  GeV as the natural scale for a more fundamental framework.

The unification of scalar masses is motivated by a similar, although more speculative, argument. Consider the mass parameter  $m_{H_u}^2$ , which, from Eq. (5), plays the critical role in determining the weak scale for all moderate and large values of  $\tan \beta$  ( $\tan \beta \gtrsim 5$ ). For theories with a universal scalar mass, it is a remarkable fact that the renormalization group trajectories of  $m_{H_u}^2$  for various initial conditions  $m_0$ , when evolved to low scales, meet with high precision at a point, the weak scale [21,22]. This focusing, which requires that the top quark mass be within  $\sim 5$  GeV of its measured value, implies that the electroweak potential is highly insensitive to  $m_0$ . The longstanding problems of supersymmetry with respect to CP violation, proton decay, etc. can therefore be ameliorated without fine-tuning, simply by assuming large scalar masses. Although the focusing property holds more generally, its minimal and most concrete realization is in theories with a universal scalar mass. The assumption of a universal (and large) scalar mass is therefore motivated by the fact that it

provides a simple and elegant solution to several well-known phenomenological problems of weak scale supersymmetry.

Given these motivations for minimal supergravity, we now consider their implications for neutralino dark matter. Gaugino mass unification implies  $M_1 \simeq M_2/2 \simeq 0.4M_{1/2}$ . Dark matter is therefore never Wino-like,<sup>1</sup> and in fact, throughout parameter space,  $|a_2|^2 < 0.07$ . Additional insights follow from re-writing Eq. (5) in terms of GUT scale parameters. For  $A_0 = 0$  and  $\tan\beta = 10$ , for example,

$$\frac{1}{2}m_Z^2 \approx -0.02 m_0^2 + 0.7 M_{1/2}^2 - \mu^2, \quad (6)$$

where the numerical coefficients of the first two terms vary fractionally by  $\mathcal{O}(10\%)$  in the  $(m_0, M_{1/2})$  plane [21,22]. The coefficient of  $m_0^2$  is highly suppressed [30,31]. This is another expression of the focusing behavior discussed above, and implies that multi-TeV values of  $m_0$  do not involve significant large fine-tuning. The coefficient is also negative. For fixed  $M_{1/2}$ , as  $m_0$  increases,  $|\mu|$  decreases, and the LSP becomes increasingly Higgsino-like. This is important, because even a 10% Higgsino admixture drastically affects the phenomenology. In Figs. 1 and 2 we show the LSP mass and gaugino fraction in the  $(m_0, M_{1/2})$  plane. For large  $m_0 \gtrsim 1$  TeV, we find a region, previously ignored, where the LSP has significant Higgsino component.

In Figs. 1 and 2, we also indicate the regions of cosmologically interesting relic densities; see Ref. [6] for details. A generous range is  $0.025 \leq \Omega_\chi h^2 \leq 1$ , where the lower bound is the requirement that neutralino dark matter explain galactic rotation curves and the upper bound follows from the lifetime of the universe. Above this shaded region,  $\Omega h^2 > 1$ ; below,  $\Omega h^2 < 0.025$ . The range  $0.1 \leq \Omega_\chi h^2 \leq 0.3$  is most preferred by current limits. Our relic density calculation is not trustworthy in the black region, where there is a Higgs scalar resonance, and very near to the left and right borders of the excluded region, where co-annihilation is important [32–35]. In the bulk of parameter space, however, these effects are negligible. For all  $\tan\beta$ , cosmologically interesting densities are possible for  $m_0 \gtrsim 1$  TeV. For  $\tan\beta = 10$ , the cosmologically preferred region contains gaugino-Higgsino dark matter.

In contrast to  $m_0$ , the parameters  $M_{1/2}$  and  $\mu$  enter Eq. (6) with  $\mathcal{O}(1)$  coefficients. Naturalness therefore requires that the LSP mass (and, in fact, the masses of all four neutralinos and both charginos) should not be too far above the electroweak scale. While in principle it is possible that in some fundamental framework  $M_{1/2}$  and  $\mu$  are correlated precisely in a way that allows both parameters to be large without fine-tuning (a possibility considered in Ref. [36]), no such framework has been found to date. Barring such a possibility, extreme values such as  $M_{1/2}, \mu \sim 50$  TeV require a fine-tuning of 1 part in  $10^6$  and destroy one of the prime motivations for weak-scale supersymmetry. We therefore regard such large values as highly disfavored, and we will focus on neutralino masses of order 100 GeV.

Neutralinos annihilate through a variety of channels. The three leading processes are shown in Fig. 3. (Note that co-annihilation, while potentially important in determining

---

<sup>1</sup>Wino-like LSPs exist in other frameworks, but typically they annihilate far too quickly to be cosmologically relevant [28]. Interesting relic densities are possible, however, if there is some mechanism of late production [29].




FIG. 1. Contours of constant LSP mass  $m_\chi$  in GeV in the  $(m_0, M_{1/2})$  plane for  $A_0 = 0$ ,  $\mu > 0$ ,  $m_t = 174$  GeV, and two representative values of  $\tan \beta$ . The green shaded regions are excluded by the requirement that the LSP be neutral (left) and by the chargino mass limit of 95 GeV (bottom and right). We have also delineated the regions with potentially interesting values of the LSP relic abundance:  $0.025 \leq \Omega_\chi h^2 \leq 1$  (yellow) and  $0.1 \leq \Omega_\chi h^2 \leq 0.3$  (light blue). In the black region,  $|2m_\chi - m_h| < 5$  GeV, and neutralino annihilation is enhanced by a Higgs resonance.




FIG. 2. Contours of constant gaugino fraction  $R_\chi$  in percent, for the same values of the parameters as in Fig. 1.

relic densities in the early universe, is negligible now.) Annihilation into gauge bosons is of particular importance, as these processes lead to more energetic and striking signals. The  $WW$  cross section relies on  $W\chi\chi_i^\pm$  interactions. The only such couplings allowed by gauge invariance are  $W\tilde{H}^0\tilde{H}^\pm$  and  $W\tilde{W}^0\tilde{W}^\pm$ . However, as noted above, gaugino mass unification implies that the Wino content of the LSP is always negligible. A large  $WW$  cross section is therefore possible only when the LSP has a significant Higgsino component. The same




FIG. 3. Three leading neutralino annihilation channels.




FIG. 4. Contours of constant  $\sigma v$  in pb for (a)  $\chi\chi \rightarrow WW$  and (b)  $\chi\chi \rightarrow ZZ$ . We fix  $A_0 = 0$ ,  $\mu > 0$ ,  $m_t = 174$  GeV, and  $\tan\beta = 10$ .

conclusion holds for the  $ZZ$  process, where the  $Z\chi\chi_i^0$  interaction is possible only through  $Z\tilde{H}^0\tilde{H}^0$  couplings. In Fig. 4, we see that the annihilation cross sections for  $\chi\chi \rightarrow WW$  and  $\chi\chi \rightarrow ZZ$  are indeed highly suppressed in regions with Bino-like LSPs, but are enhanced by three to four orders of magnitude in regions with mixed gaugino-Higgsino dark matter. As we will see, this region, favored by low energy constraints, will be the most promising for all indirect signals.

Before closing this section, we note several features of Figs. 1 and 2 that will also apply to many of the following figures. Unless otherwise noted, we present results for  $A_0 = 0$ ,  $\mu > 0$ ,  $m_t = 174$  GeV, and representative values of  $\tan\beta$  as indicated.  $A_0$  governs the left-right mixing of scalars, and does not enter the neutralino sector. It is therefore largely irrelevant, especially in the regions of parameter space with observable signals — where, as we will see, the scalars are heavy and decoupled. (Besides, the most important trilinear coupling,  $A_t$ , has a weak-scale fixed point, and so is only weakly sensitive to  $A_0$ .) Our dark matter results are rather insensitive to the sign of  $\mu$ , but the choice  $\mu > 0$  is motivated by the constraint from  $B \rightarrow X_s \gamma$ . Finally, perturbativity of Yukawa couplings limits  $\tan\beta$  to the range  $1 \lesssim \tan\beta \lesssim 60$ . Low values of  $\tan\beta \lesssim 3$  are now being excluded by the LEP Higgs search. In the remaining interval, models with moderate and high values may have qualitatively different behavior, as processes proportional to down-type Yukawa couplings are enhanced by  $\tan\beta$ . We therefore typically present results for two representative values,



one in each range. Plots of many other quantities, including all physical slepton, squark, and Higgs masses in the  $(m_0, M_{1/2})$  plane, including the high  $m_0$  region, may be found in Refs. [21–23].

Finally, in the following three sections, we will *not* include the effects of variations in thermal relic density in our signal rates, but rather assume, for concreteness, a fixed local neutralino density. The results are then more transparent, and are applicable to general scenarios, such as those in which a late source of neutralino production is present. Of course, in the simplest scenario, models with  $\Omega_\chi h^2 > 1$  are excluded, and those with an underabundance of neutralinos are disfavored and imply suppressed or negligible dark matter signals. This should be kept in mind in the following sections. In Secs. VI and VII, we will combine all these considerations, and focus on the most preferred regions.

### III. NEUTRINOS

When neutralinos pass through astrophysical objects, they may be slowed below escape velocity by elastic scattering. Once captured, they then settle to the center, where their densities and annihilation rates are greatly enhanced. While most of their annihilation products are immediately absorbed, neutrinos are not. High energy neutrinos from the cores of the Earth [37–40] and Sun [39,41–46] are therefore promising signals for indirect dark matter detection.

The formalism for calculating neutrino fluxes from dark matter annihilation is well developed. (See Ref. [47] for a review.) The neutrino flux depends first and foremost on the neutralino density, which is governed by the competing processes of gravitational capture and neutralino annihilation. If  $N$  is the number of neutralinos in the Earth or Sun,  $\dot{N} = C - AN^2$ , where  $C$  is the capture rate and  $A$  is the total annihilation cross section times relative velocity per volume. The present neutralino annihilation rate is then

$$\Gamma_A = \frac{1}{2}AN^2 = \frac{1}{2}C \tanh^2(\sqrt{CA}t_\odot), \quad (7)$$

where  $t_\odot \approx 4.5$  Gyr is the age of the solar system.

Captured neutralinos then annihilate through the processes of Fig. (3). As  $\chi\chi \rightarrow f\bar{f}$  is helicity-suppressed, neutrinos are produced only in the decays of primary annihilation products. Typical neutrino energies are then  $E_\nu \sim \frac{1}{2}m_\chi$  to  $\frac{1}{3}m_\chi$ , with the most energetic spectra resulting from  $WW$ ,  $ZZ$ , and, to a lesser extent,  $\tau\bar{\tau}$ . After propagating to the Earth’s surface, neutrinos are detected through their charged-current interactions. The most promising signal is from upward-going muon neutrinos that convert to muons in the surrounding rock, water, or ice, producing through-going muons in detectors. The detection rate for such neutrinos is greatly enhanced for high energy neutrinos, as both the charged-current cross section and the muon range are proportional to  $E_\nu$ .

The calculation of muon fluxes from neutralino annihilation in the Earth and Sun is on reasonably firm footing, as it depends only on the local dark matter density and is insensitive to details of halo modeling.<sup>2</sup> Nevertheless, the calculation is involved, primarily as a result of

---

<sup>2</sup>It has been suggested that muon fluxes may be enhanced, by up to two orders of magnitude,




FIG. 5. The filling parameter  $\sqrt{CA} t_\odot$  for the Earth. We assume neutralino velocity dispersion  $\bar{v} = 270$  km/s and a local density of  $\rho_0 = 0.3$  GeV/cm<sup>3</sup>.

complications in evaluating capture rates [50–52] and, in the case of the Sun, in propagating the neutrinos from the core to the surface [39,53,54]. Here we make use of the procedure of Refs. [47,55]. For other analyses, see Refs. [56–58,17], as well as those motivated by the Tevatron  $e^+e^-\gamma\gamma$  event [59,60] and by the DAMA annual modulation signal [61,62].




FIG. 6. As in Fig. 5, but for the Sun.

The ‘filling parameters’  $\sqrt{CA} t_\odot$  for the Earth and Sun are given in Figs. 5 and 6. For the

---

due to capture of neutralinos in highly eccentric solar system orbits [48]. The magnitude of the enhancement depends on details of the neutralino parameters and involves astrophysical issues still under debate [49]. We have not included it here.




FIG. 7. The capture rate of neutralinos  $C$  in  $\text{s}^{-1}$  in the center of the Earth (for  $\bar{v} = 270$  km/s and  $\rho_0 = 0.3$   $\text{GeV}/\text{cm}^3$ ).

Sun,  $\sqrt{CA}t_\odot \gg 1$  for all supersymmetry parameters. The neutralino density has therefore reached equilibrium, and the annihilation rate is at full strength, with  $\Gamma_A \approx C/2$ . For the Earth, however, typically  $\sqrt{CA}t_\oplus \ll 1$ , and the annihilation rate is  $\Gamma_A \approx \frac{1}{2}C^2 A t_\oplus^2$  and far from maximal. As we will see, this plays an important part in reducing the Earth's signal below the Sun's.




FIG. 8. Same as in Fig. 7, but for the Sun.

The other major ingredient in the muon flux computation is the estimate of the neutralino capture rate  $C$ , which is shown in Figs. 7 and 8 for the Earth and Sun, respectively. The elemental compositions of the Earth and Sun are given in Ref. [11]. A quick comparison of Figs. 7 and 8 reveals that, not surprisingly, a large astrophysical body like the Sun is much more efficient in trapping neutralinos. The Earth's capture rate is, however, enhanced by the




FIG. 9. Muon flux from the Earth in  $\text{km}^{-2} \text{yr}^{-1}$  (for  $\bar{v} = 270 \text{ km/s}$  and  $\rho_0 = 0.3 \text{ GeV/cm}^3$ ).

iron resonance for very light neutralino masses  $m_\chi \sim 50 \text{ GeV}$ . The  $\tan \beta$  dependence is also noteworthy. The capture rate in the Earth is determined primarily by the spin-independent elastic scattering cross section for  $\chi q \rightarrow \chi q$  through  $s$ -channel squarks and  $t$ -channel Higgs boson exchange. All amplitudes require chirality flips, either through Higgs interactions, squark mass insertions, or quark mass insertions. The first two are proportional to  $\tan \beta$  and therefore dominate for moderate and large  $\tan \beta$ , leading to  $C \sim \tan^2 \beta$ . In contrast, for the Sun, the  $\tan \beta$  dependence is minimal; the dominant contribution is from axial-vector scattering off Hydrogen, which is largely independent of  $\tan \beta$ .




FIG. 10. As in Fig. 9, but for the Sun.

Muon flux rates from the Earth and Sun are presented in Figs. 9 and 10. Consistent with previous studies, we find that the flux rate is indeed small in regions of parameter space with  $m_0 < 1 \text{ TeV}$  and Bino-like LSPs. However, for  $m_0 > 1 \text{ TeV}$ , in the region where  $m_\chi > m_W$

TABLE I. Current and planned neutrino experiments. We list also each experiment's (expected) start date, physical dimensions (or approximate effective area), muon threshold energy  $E_\mu^{\text{thr}}$  in GeV, and 90% CL flux limits for the Earth  $\Phi_\mu^\oplus$  and Sun  $\Phi_\mu^\odot$  in  $\text{km}^{-2} \text{yr}^{-1}$  for half-cone angle  $\theta \approx 15^\circ$  when available.

Experiment	Type	Date	Dimensions	$E_\mu^{\text{thr}}$	$\Phi_\mu^\oplus$	$\Phi_\mu^\odot$
Baksan [64]	Underground	1978	$17 \times 17 \times 11 \text{ m}^3$	1	$6.6 \times 10^3$	$7.6 \times 10^3$
Kamiokande [65]	Underground	1983	$\sim 150 \text{ m}^2$	3	$10 \times 10^3$	$17 \times 10^3$
MACRO [66]	Underground	1989	$12 \times 77 \times 9 \text{ m}^3$	1.5 *	$3.2 \times 10^3$	$6.5 \times 10^3$
Super-Kamiokande [67]	Underground	1996	$\sim 1200 \text{ m}^2$	1.6	$1.9 \times 10^3$	$5.0 \times 10^3$
Baikal NT-96 [68]	Underwater	1996	$\sim 1000 \text{ m}^2$	10	$15 \times 10^3$	
AMANDA B-10 [69]	Under-ice	1997	$\sim 1000 \text{ m}^2^\dagger$	$\sim 25$	$44 \times 10^3^\dagger$	
Baikal NT-200 [68]	Underwater	1998	$\sim 2000 \text{ m}^2$	$\sim 10$		
AMANDA II [70]	Under-ice	2000	$\sim 3 \times 10^4 \text{ m}^2$	$\sim 50$		
NESTOR <sup>§</sup> [71]	Underwater	2000	$\sim 10^4 \text{ m}^2^\ddagger$	few		
ANTARES [72]	Underwater	2003	$\sim 2 \times 10^4 \text{ m}^2^\ddagger$	$\sim 5\text{--}10$		
IceCube [70]	Under-ice	2003-8	$\sim 10^6 \text{ m}^2$			

\* 2 GeV for Sun.  $^\dagger$  Hard spectrum,  $m_\chi = 100 \text{ GeV}$ .  $^\S$  One tower.  $^\ddagger$   $E_\mu \sim 100 \text{ GeV}$ .

and the dark matter is a gaugino-Higgsino mixture, the fluxes are greatly enhanced. Here, annihilation to gauge bosons is unsuppressed, resulting in a hard neutrino spectrum and large muon fluxes. In this region, the rates from the Sun are large for all values of  $\tan \beta$ . For the Earth, we see that, despite the close proximity of the Earth's center, the muon fluxes are typically suppressed by several orders of magnitude relative to those from the Sun. However, reasonably large rates are possible even for the Earth for large  $\tan \beta$ , and particularly for very light neutralinos, where the capture rate is enhanced by the iron resonance, as discussed above.

The theoretical predictions of Figs. 9 and 10 can be compared with the experimental sensitivities of ongoing and near future detectors [63]. These experiments, along with their more salient characteristics and flux limits (where available), are listed in Table I. The flux limits depend on the expected angular dispersion in the signal. This dispersion has two possible origins. One is the source: although neutralinos from the Sun are essentially a point source, in the Earth, 98% of neutralino annihilations occur within a cone of half-angle  $8.6^\circ \sqrt{50 \text{ GeV}/m_\chi}$  [51,52]. The second is the angle  $\theta_{\text{rms}} \approx 13^\circ \sqrt{25 \text{ GeV}/E_\nu}$  between the neutrino and its daughter muon. As  $E_\nu \lesssim m_\chi/2$ ,  $\theta_{\text{rms}}$  is typically the dominant effect. The flux limits listed are for half-cone sizes of  $15^\circ$ , corresponding roughly to  $m_\chi \sim 50 \text{ GeV}$ . Heavier neutralinos will produce more collimated muons, allowing smaller cone sizes with reduced backgrounds. The improved limits for smaller cone sizes may be found in the references.

Comparing Figs. 9 and 10 with Table I, we find that present limits do not significantly constrain the minimal supergravity parameter space. However, given that the effective area of neutrino telescope experiments is expected to increase by 10 to 100 in the next few years, muon fluxes of order  $10\text{--}100 \text{ km}^{-2} \text{yr}^{-1}$  may be within reach. Such sensitivities are typically not sufficient to discover Bino-like LSPs, unless they are light and  $\tan \beta$  is large. But they

have an excellent opportunity to detect dark matter in the mixed gaugino-Higgsino dark matter scenarios, which, as we have emphasized above, are preferred by low energy particle physics constraints.

Muon energy thresholds, listed in Table I, are not included in Figs. 9 and 10. Since the muon detection rate is dominated by high energy muons as noted above, the threshold energy is typically not important, especially in the regions where a detectable signal is expected. This is not the case for all detectors, however. For example, since muons lose 0.26 GeV per meter in water and ice, neutrino telescopes requiring track lengths of  $\sim 100$  m will have thresholds of order  $\sim 25$  GeV. The dependence on threshold energy has been studied in Refs. [73,18], where it was found that for threshold energies of  $E_\mu^{\text{thr}} \sim \frac{1}{4}m_\chi$  to  $\frac{1}{6}m_\chi$ , the loss of signal is substantial. Low threshold energies in neutrino telescopes are clearly very important for dark matter detection. This conclusion is further strengthened by considerations of naturalness, which favor low neutralino masses.

#### IV. PHOTONS

High-energy photons provide a unique signal of dark matter annihilation. They point back to their source, and their energy distribution is directly measurable, at least in principle. For these reasons, given sufficient angular and energy resolution in gamma ray detectors, a variety of signals may be considered.

The photon signal may arise from the galactic center [74–76], the galactic halo [77,78], or even from extra-galactic sources [20]. We will consider the galactic center, where large enhancements in dark matter density are possible [19,79]. In contrast to the neutrino signal considered in Sec. III, the photon flux is highly sensitive to halo model parameters. Fortunately, the problem may be separated into two parts: one containing all halo model dependence, and the other all particle physics uncertainties. Given the predicted photon fluxes for a reference halo model, the predictions for all other halo models are then easily determined.

The photon energy distribution receives two types of contributions: line and continuum. The former results from the loop-mediated processes  $\chi\chi \rightarrow \gamma\gamma$  [80,81] and  $\chi\chi \rightarrow \gamma Z$  [82]. Because dark matter in the halo is extremely non-relativistic, photons from these processes have an energy width of only  $\Delta E_\gamma/E_\gamma \sim 10^{-3}$  and are effectively mono-energetic. While this signal would be the most spectacular of all possible indirect signals, its rates are, of course, suppressed [83]. In a model-independent survey, Bergström, Ullio, and Buckley [19] have found that the photon line may be observable for neutralinos with a large Higgsino component, assuming a cuspy halo profile, such as that of Navarro, Frenk, and White [84], and telescopes with small angular acceptances  $\sim 10^{-5}$  sr.

On the other hand, photons may also be produced in the cascade decays of other primary annihilation products. In contrast to the line signal, cascade decays produce a large flux of photons with a continuum of energies. This signal is far less distinctive and will almost certainly require additional confirmation to unambiguously distinguish it from background or other exotic sources. Nevertheless, we will focus here on the continuum signal, as it will provide the first hint of dark matter from gamma ray astronomy.

The differential photon flux along a direction that forms an angle  $\psi$  with respect to the direction of the galactic center is

$$\frac{d\Phi_\gamma}{d\Omega dE} = \sum_i \frac{dN_\gamma^i}{dE} \sigma_{iv} \frac{1}{4\pi m_\chi^2} \int_\psi \rho^2 dl , \quad (8)$$

where the sum is over all annihilation channels and  $\rho$  is the neutralino mass density. All of the halo model dependence is isolated in the integral, which, following Ref. [19], we write in the dimensionless form

$$J(\psi) = \frac{1}{8.5 \text{ kpc}} \left( \frac{1}{0.3 \text{ GeV/cm}^3} \right)^2 \int_\psi \rho^2 dl . \quad (9)$$

The integral is along the line of sight. Assuming a spherical halo, the mass density is given by  $\rho = \rho(r)$ , where  $r^2 = l^2 + R_0^2 - 2lR_0 \cos \psi$ , and  $R_0 \approx 8.5 \text{ kpc}$  is the solar distance to the galactic center.

The photon flux is, of course, maximized for  $\psi = 0$ , but it must be averaged over the field of view. The result is

$$\Phi_\gamma(E_{\text{thr}}) = 5.6 \times 10^{-10} \text{ cm}^{-2} \text{ s}^{-1} \times \sum_i \int_{E_{\text{thr}}}^{m_\chi} dE \frac{dN_\gamma^i}{dE} \left( \frac{\sigma_{iv}}{\text{pb}} \right) \left( \frac{100 \text{ GeV}}{m_\chi} \right)^2 \bar{J}(\Delta\Omega) \Delta\Omega , \quad (10)$$

where

$$\bar{J}(\Delta\Omega) \equiv \frac{1}{\Delta\Omega} \int_{\Delta\Omega} J(\psi) d\Omega , \quad (11)$$

and  $\Delta\Omega$  is the solid angle of the field of view centered on  $\psi = 0$ .  $E_{\text{thr}}$  is the lower threshold energy; detectors also have upper cutoffs, but these are typically irrelevant, as the energy distribution falls steeply with energy.  $\bar{J}$  has been studied for a variety of halo models in Ref. [19]. For a typical atmospheric Cherenkov telescope (ACT) acceptance of  $\Delta\Omega = 10^{-3} \text{ sr}$ , the modified isothermal profile described by  $\rho(r) \propto [1 + (r/a)^2]^{-1}$  yields  $3 \lesssim \bar{J} \lesssim 10^3$ . On the other hand, cuspy halos lead to values of  $\bar{J}$  as large as  $10^5$ . (Such singular profiles have recently been argued to be incompatible with neutralino dark matter, however, based on radio emission from neutralino annihilation near the black hole at the galactic center [85].) We will choose a moderate reference value  $\bar{J}(10^{-3}) = 500$ , which is within the allowed ranges of both the modified isothermal and cuspy halos. The factorizability of the photon flux implies that our results can be scaled to all other halo models easily.

The particle physics model dependence enters through all the other factors of Eq. (8). The energy integral is roughly  $\int dE dN_\gamma^i/dE \sim 0.5$  for all  $i$ , but the energy distribution depends significantly on the annihilation channel. The differential gamma ray multiplicity has been simulated for light and heavy neutralinos in Refs. [86] and [19], respectively. The spectrum for the most important annihilation channels is described well by  $dN_\gamma/dx = a e^{-bx}/x^{1.5}$ , where  $x \equiv E_\gamma/m_\chi$  and  $(a, b) = (0.73, 7.76)$  for  $WW$  and  $ZZ$  [19],  $(1.0, 10.7)$  for  $b\bar{b}$ ,  $(1.1, 15.1)$  for  $t\bar{t}$ , and  $(0.95, 6.5)$  for  $u\bar{u}$ . We neglect Higgs boson final states, as they never have branching fraction greater than 7%. For the  $gg$  final state, we use the light quark distribution. Our crude approximation for gluons is relevant only in isolated regions with Bino-like LSPs where, as we will see, the signal is unobservable. With the exception of the irrelevant light quark distribution, the  $WW$  and  $ZZ$  distributions produce the most energetic photons.

The photon flux  $\Phi_\gamma(E_{\text{thr}})$  is given in Figs. 11–13 for threshold energies of 1, 10 and 50 GeV. The maximal rates are found in the region of parameter space with mixed gaugino-Higgsino dark matter, and are insensitive to  $\tan\beta$ . Here branching ratios to gauge bosons




FIG. 11. Photon flux  $\Phi_\gamma(E_{\text{thr}})$  in  $\text{cm}^{-2} \text{s}^{-1}$  from a  $10^{-3}$  sr cone centered on the galactic center for a threshold energy of  $E_{\text{thr}} = 1$  GeV. We assume halo model parameter  $\bar{J} = 500$ . Results for other halo models may be obtained by scaling to the appropriate  $\bar{J}$  (see text).




FIG. 12. As in Fig. 11, but for the photon energy threshold  $E_{\text{thr}} = 10$  GeV.




FIG. 13. As in Fig. 11, but for the photon energy threshold  $E_{\text{thr}} = 50$  GeV.



TABLE II. Some of the current and planned  $\gamma$  ray detector experiments with sensitivity to photon energies  $10 \text{ GeV} \lesssim E_\gamma \lesssim 300 \text{ GeV}$ . We list each experiment's (proposed) start date and expected  $E_\gamma$  coverage in GeV. The energy ranges are approximate. For experiments constructed in stages, the listed threshold energies will not be realized initially. See the references for details.

Experiment	Type	Date	$E_\gamma$ Range
EGRET [87]	Satellite	1991-2000	0.02–30
STACEE [88]	ACT array	1998	20–300
CELESTE [89]	ACT array	1998	20–300
ARGO-YBJ [90]	Air shower	2001	100–2,000
MAGIC [91]	ACT	2001	10–1000
AGILE [92]	Satellite	2002	0.03–50
HESS [93]	ACT array	2002	40–5000
AMS/ $\gamma$ [94]	Space station	2003	0.3–100
CANGAROO III [95]	ACT array	2004	30–50,000
VERITAS [96]	ACT array	2005	50–50,000
GLAST [97]	Satellite	2005	0.1–300

are large, and the photon spectrum hard. In the rest of parameter space,  $b\bar{b}$  is an important final state, and  $\Phi_\gamma(E_{\text{thr}})$  is enhanced by  $\tan \beta$ .

In the past, the high energy photon spectrum with  $10 \text{ GeV} \lesssim E_\gamma \lesssim 300 \text{ GeV}$  has been largely unexplored. Ground-based detectors, such as the Whipple 10m telescope, have large effective areas, but have traditionally been limited to energies above  $\sim 300 \text{ GeV}$ . Space-based detectors, such as EGRET, have been sensitive to photon energies up to  $\sim 20 \text{ GeV}$ , but are limited above this energy by their small effective area. There has therefore been an unexplored gap at intermediate energies, which happens to overlap substantially with the range of energies most favored by supersymmetric dark matter.

The experimental situation is changing rapidly, however. Currently, two heliostat arrays, STACEE and CELESTE, are running with sensitivity in the range  $20 \text{ GeV} \lesssim E_\gamma \lesssim 300 \text{ GeV}$ , and many more experiments with greatly improved sensitivity are expected in the next few years. Upcoming experiments with sensitivity to  $\gamma$  rays with  $10 \text{ GeV} \lesssim E_\gamma \lesssim 300 \text{ GeV}$  are listed in Table II.

An important figure of merit for the detection of  $\gamma$  rays from the galactic center is the point source flux sensitivity. A compilation of previous estimates of flux sensitivities is given in Fig. 14 for EGRET, STACEE, CELESTE, ARGO-YBJ, MAGIC, AGILE, HESS [93], AMS/ $\gamma$  [94], CANGAROO III [95], VERITAS [96], and GLAST [97]. The flux sensitivities for the first six experiments are from Ref. [92], and those for the remaining experiments are from the references listed. The point flux sensitivities are, of course, dependent on the source's location and energy spectrum. They are also subject to a variety of other experimental uncertainties and assumptions; see the references for details. A typical estimate [96] assumes background extrapolated from EGRET data [87], and a signal distribution  $dN_\gamma/dE \propto E^{-2.5}$ . Detector efficiencies and cuts are included, and a  $5\sigma$  signal with at least 10 photons is required. 50 hours of observation is assumed for telescopes, and one year of an all sky survey for the space-based detectors. The arrow for AMS/ $\gamma$  in Fig. 14 indicates that a published estimate exists only for  $E_{\text{thr}} = 1 \text{ GeV}$ , but flux sensitivity at some level




FIG. 14. Photon fluxes  $\Phi_\gamma(E_{\text{thr}})$  as a function of threshold energy  $E_{\text{thr}}$  for  $A_0 = 0$ ,  $\mu > 0$ ,  $m_t = 174$  GeV, and halo parameter  $\bar{J} = 500$ . The four models have relic density  $\Omega_\chi h^2 \approx 0.15$ , and are specified by  $(\tan \beta, m_0, M_{1/2}, m_\chi, R_\chi) = (10, 100, 170, 61, 0.93)$  (dotted),  $(10, 1600, 270, 97, 0.77)$  (dashed),  $(10, 2100, 500, 202, 0.88)$  (dot-dashed), and  $(50, 1000, 300, 120, 0.96)$  (solid), where all masses are in GeV. Point source flux sensitivity estimates for several gamma ray detectors are also shown. (Care should be taken in comparing these sensitivities to the predicted fluxes — see text.)

can be expected out to the detector limit of 100 GeV.

The experimental sensitivities presented clearly cannot be interpreted as future dark matter discovery contours. The neutralino signal has a different energy spectrum than assumed, and the background in the direction of the galactic center is larger, due to the diffuse  $\gamma$  ray emission from the galactic disk [98], which enhances  $\sqrt{B}$  by a factor of  $\sim 5$  [99]. (This last fact implies that for some halo profiles, it may be advantageous to center the field of view away from the galactic center. This optimization may significantly reduce potential losses in signal significance.) In addition, there are many ambiguities in background calibration, and, as noted above, the continuum signal is not sufficiently distinct for a simple excess to identify the source as dark matter annihilation. However, the flux sensitivities of Fig. 14 do clearly portend substantial progress in the next few years, and can serve as rough indications of what signal levels will be detectable.

The expected fluxes for four supersymmetry models and  $\bar{J} = 500$  are also shown in Fig. 14. Although there is a large uncertainty from halo model dependence, it is clear that detectable signals are possible. At present, EGRET data is not overly constraining, although halo models with large  $\bar{J} \sim 5000$  are within EGRET sensitivity and may even explain a flattening of the spectrum. In the future, AMS/ $\gamma$  and AGILE will improve this sensitivity, and MAGIC, HESS, and CANGAROO III may see excesses for  $\bar{J} \sim 500$ . Finally, GLAST will provide the greatest sensitivity of all, probing halo models with  $\bar{J}$  as low as  $\bar{J} \sim 50$ .

If a significant excess is found in future experiments, its dark matter origin can be tested in a variety of ways. Confirmation from other searches for dark matter or supersymmetry would be the most satisfying possibility. As we will see in Sec. VI, the neutrino and positron signals probe similar models, so a coincidence of various signals is a distinct possibility. However, it has also been suggested that an angular distribution of photons that does not follow the galactic disk and bulge may be a powerful diagnostic [19]. Also, as the dark matter signal has a shape differing from the background, detailed likelihood fits to the photon energy distribution may also be a useful tool, although far beyond the scope of this work. It seems clear, in any case, that for reasonable halo models and supersymmetry parameters, meaningful  $\gamma$  ray signals in the next few years are possible, particularly with gaugino-Higgsino dark matter.

## V. POSITRONS

An excess of cosmic anti-particles and anti-matter from dark matter annihilation may be detected in space-based or balloon-borne experiments. The positron signal is perhaps the most promising [100–104]. In the past, soft anti-protons with energies  $\sim 100$  MeV have also been considered [105,106]. However, recent work finds larger background than previously expected, complicating the identification of a anti-proton signal [107,108]. Anti-deuterium has also been suggested as a possibility [109].

The positron background is most likely to be composed of secondaries produced in the interactions of cosmic ray nuclei with interstellar gas, and is expected to fall as  $\sim E_{e^+}^{-3.1}$ . At energies below 10 GeV, however, this background is subject to large uncertainties from the effects of the solar wind [103,104]. The soft positron spectrum also varies depending on the orbit path of the experiment. At high energies, these effects are negligible. In addition, positrons lose energy through a variety of processes, and so hard ones must typically be produced within a few kpc [103,104]. For this reason, the hard spectrum is relatively insensitive to variations in the halo profile near the galactic center. The dark matter signal is therefore most promising at high energies, where the background is relatively small and well understood.

The differential positron flux is [104]

$$\frac{d\Phi_{e^+}}{d\Omega dE} = \frac{\rho_0^2}{m_\chi^2} \sum_i \sigma_i v B_{e^+}^i \int dE_0 f_i(E_0) G(E_0, E) , \quad (12)$$

where  $\rho_0$  is the local neutralino mass density, the sum is over all annihilation channels, and  $B_{e^+}^i$  is the branching fraction to positrons in channel  $i$ . The source function  $f(E_0)$  gives the initial positron energy distribution from neutralino annihilation.  $G(E_0, E)$  is the Green's function describing positron propagation in the galaxy [110] and contains all the halo model dependence.

For the reasons mentioned above, processes yielding hard positrons are by far the most important for dark matter discovery. The ‘positron line’ signal from  $e^+e^-$  is helicity-suppressed. It may be enhanced, for example, in the case of Bino-like LSPs if selectrons are much lighter than all other scalars, but this possibility is highly unmotivated. To an excellent approximation, then, hard positrons arise from  $\chi\chi \rightarrow WW, ZZ$ , followed by the direct

decay of gauge bosons to positrons. Assuming unpolarized gauge bosons,  $f$  is the familiar flat distribution with endpoints determined by the gauge boson and neutralino masses. The Green's function  $G$  has been modeled by Moskalenko and Strong in Ref. [104] in a framework that consistently reproduces a wide range of observational data from anti-protons, nuclei, electrons, positrons, and photons.

Combining all of these results, the differential positron flux may be written as

$$E^2 \frac{d\Phi_{e^+}}{d\Omega dE} = 0.027 \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ GeV} \times \left( \frac{\rho_0}{0.3 \text{ GeV/cm}^3} \right)^2 \left( \frac{100 \text{ GeV}}{m_\chi} \right)^2 \sum_i \frac{\sigma_i v}{\text{pb} \cdot \beta_i} B_{e^+}^i \int_{z_-^i}^{z_+^i} dz g(z, E/m_\chi), \quad (13)$$

where

$$\beta_{WW,ZZ} = \left( 1 - m_{W,Z}^2/m_\chi^2 \right)^{1/2}, \quad (14)$$

$$z_\pm^i = (1 \pm \beta_i)/2, \quad (15)$$

$$B_{e^+}^{WW} = B(W^+ \rightarrow e^+ \nu) = 0.11, \quad (16)$$

$$B_{e^+}^{ZZ} = 2B(Z \rightarrow e^+ e^-) = 0.067, \quad (17)$$

and the reduced Green's function is

$$g(z, E/m_\chi) \equiv 10^{a \log_{10}^2 E + b \log_{10} E + c} \theta(z - E/m_\chi) + 10^{w \log_{10}^2 E + x \log_{10} E + y} \theta(E/m_\chi - z), \quad (18)$$

where  $E$  is in GeV and the ( $z$ -dependent) coefficients  $a, b, c$  and  $w, x, y$  are tabulated in Ref. [104] for different halo profiles. As mentioned above, at high energies, these coefficients are fairly independent of the halo model, as high energy positrons originate in our solar neighborhood, where all profiles give similar densities. We adopt coefficients corresponding to the modified isothermal distribution with halo size 4 kpc. For large  $m_\chi$ , the integral of Eq. (13) is insensitive to  $m_\chi$ , and so the differential positron flux scales as  $\sim 1/m_\chi^4$ . Neutralinos with mass not far above  $m_W$  are therefore most easily detected.

In Fig. 15, we show three sample spectra for supersymmetry models yielding relic abundances of  $\Omega_\chi h^2 \approx 0.15$ . Two background spectra from Ref. [104] are also shown. The signal rates are significantly suppressed relative to those of Refs. [102,104], where the dark matter was assumed to be Higgsino-like. Higgsino-like dark matter is highly disfavored, however, as, unless it is unnaturally heavy, it annihilates too strongly to leave interesting relic abundances. As is evident from Fig. 2, in the allowed minimal supergravity parameter space the LSP is far from pure Higgsino-like, particularly in the region with preferred relic density.

As positrons result from two-body decay, we expect the signal, and the signal to background ratio  $S/B$ , to be peaked near  $m_\chi/2$ . This is evident in the three examples give in Fig. 15. In Fig. 16, we plot the optimal energy  $E_{\text{opt}}$  at which the signal to background ratio is maximized. Our fit to background C is  $E^2 d\Phi_{e^+}/d\Omega dE = 1.16 \times 10^{-3} E^{-1.23}$ , where  $E$  is in GeV. Comparing with Fig. 1, we see that  $E_{\text{opt}}$  is indeed approximately  $m_\chi/2$  throughout parameter space. In Fig. 17, we plot  $S/B$  at  $E_{\text{opt}}$ .  $S/B$  is substantial only in the gaugino-Higgsino region with  $m_\chi > m_W$ .

Figs. 16 and 17 imply that the best experimental hope for indirect detection of dark matter through positrons is in experiments sensitive to positron energies above  $\sim 50$  GeV.




FIG. 15. The differential positron flux for three of the four sample points in Fig. 14. The curves labeled C and HEMN are background models from Ref. [104]; the dotted curve is our fit to C.




FIG. 16. The optimal positron energy  $E_{\text{opt}}$  in GeV at which the signal to background ratio  $S/B$  is maximized.

In the next two to three years, both PAMELA, a satellite detector, and AMS-02, an experiment to be placed on the International Space Station, will satisfy this requirement. These experiments and other recently completed experiments are listed in Table III.

The expected number of positrons per GeV are listed in Table III at positron energies of 50 and 100 GeV. After integrating over some appropriate energy bin size, we see that the expected statistical errors are roughly  $\sim 10\%$  for PAMELA, and  $\sim 1\%$  for AMS-02. Of course, the signal will also be degraded by systematic errors, particularly in the background calculation. It seems likely, however, that the characteristic peaking of the dark matter signal over an interval of  $\mathcal{O}(10 \text{ GeV})$  will be distinctive. In addition, some systematic errors may be eliminated by considering the ratio  $e^+/(e^- + e^+)$ . While the positron signal is




FIG. 17. The positron signal to background ratio  $S/B$  at  $E_{\text{opt}}$ .

TABLE III. Recent and planned  $e^+$  detector experiments. We list each experiment's (expected) start date, duration, geometrical acceptance in  $\text{cm}^2 \text{ sr}$ , maximal  $E_{e^+}$  sensitivity in GeV, and (expected) total number of  $e^+$  detected per GeV at  $E_{e^+} = 50$  and 100 GeV.

Experiment	Type	Date	Duration	Acceptance	$E_{e^+}^{\text{max}}$	$\frac{dN}{dE}(50)$	$\frac{dN}{dE}(100)$
HEAT94/95 [111]	Balloon	1994/95	29/26 hr	495	50	—	—
CAPRICE94/98 [112]	Balloon	1994/98	18/21 hr	163	10/30	—	—
PAMELA [113]	Satellite	2002-5	3 yr	20	200	7	0.7
AMS-02 [114]	Space station	2003-6	3 yr	6500	1000	2300	250

typically too small for Bino-like LSPs, an excess of  $\sim 1\%$  is possible for gaugino-Higgsino dark matter. The region of detectable positron signals may be extended, however, if, for example, the halo is clumpy, or if the local density is larger than our reference value of  $0.3 \text{ GeV/cm}^3$  [103].

## VI. COMPARISON WITH OTHER SEARCHES

In the previous three sections, we have examined several indirect signals of neutralino dark matter. As emphasized in Sec. I, however, supersymmetric dark matter cannot exist in isolation, and there are many other avenues for probing supersymmetric models. We now discuss several other promising probes and their projected sensitivities, and we then compare their reaches.

Most closely linked to indirect searches are searches for dark matter scattering off nuclei in low-background detectors. The DAMA collaboration has reported evidence for an annual modulation signal [115], and the activity in this field will intensify tremendously in the next few years. (For a recent review, see, *e.g.*, Ref. [116].) Here we will use our previous results [6] to estimate the sensitivities of the direct searches. We choose CDMS (Soudan) [117] and

CRESST [118] as examples of near-term future experiments. Their projected sensitivities in neutralino-proton cross section are both of order  $\sigma_{\text{proton}} \sim 10^{-8}$  pb for  $50 \text{ GeV} \lesssim m_\chi \lesssim 500 \text{ GeV}$ . More precisely, we parameterize their sensitivities as

$$\sigma_{\text{proton}} \simeq \exp \left\{ a + b \left( \frac{m_\chi}{100 \text{ GeV}} \right) + c \left( \frac{m_\chi}{100 \text{ GeV}} \right)^2 \right\} \text{ pb} , \quad (19)$$

with  $(a, b, c) = (-17, -4.5, 3.1)$  for  $m_\chi < 84 \text{ GeV}$  and  $(-19, 0.68, -0.057)$  for  $m_\chi > 84 \text{ GeV}$ . This limit may be improved by an order of magnitude by the recently proposed GENIUS project [119], or even by CRESST itself, assuming three years of operation with improved background rejection [118].

Among high energy colliders, the Large Hadron Collider (LHC) at CERN is the ultimate supersymmetry discovery machine and will discover at least some superpartners in all of the regions of parameter space we have plotted. The LHC is scheduled to begin operation in its low luminosity mode in 2006. Before that, however, both the LEP II and Tevatron colliders have a chance to discover superpartners. The most stringent constraint from LEP II on minimal supergravity comes from chargino searches. LEP II is now concluding its run, and by the end of 2000 will improve the current chargino mass limits by about 5 GeV. If no signal is seen, this will marginally extend the bottom and right excluded regions of our figures.

The Tevatron will begin operation early in 2001. In the first two years, Run IIa will provide an integrated luminosity of  $2 \text{ fb}^{-1}$  for each detector before a temporary shutdown for a year of detector maintenance and upgrades. In the subsequent Run IIb, the data acquisition rate is expected to be about  $5 \text{ fb}^{-1}$  per year per detector. Hence, by 2006 we expect  $10 - 12 \text{ fb}^{-1}$  for each Tevatron collaboration. The Tevatron supersymmetry reach has been extensively studied recently [120], with the conclusion that there is some sensitivity, but in rather limited regions of parameter space. The most effective signal is in the clean trilepton channel [121–124] resulting from chargino-neutralino pair production, followed closely by the jets plus  $\cancel{E}_T$  channel [125] and the dileptons plus  $\tau$  jet channel [126]. The maximal reach in chargino mass is 170 GeV in the Bino LSP region at very low  $m_0$ , where the leptonic branching ratios of the electroweak gauginos are enhanced by light sleptons. This degrades rapidly at higher  $m_0$ , where hadronic decays are prominent. It also requires moderate  $\tan \beta$ . At large values of  $\tan \beta$ , decays to  $\tau$  leptons dominate the small  $m_0$  region and signatures with  $\tau$  jets must be used [125,126].

At present there are no dedicated Tevatron studies in the focus point region. (For an LHC study, see Ref. [127].) There are several important modifications to collider signals for  $m_0 > 1 \text{ TeV}$ . For example, the lighter chargino and neutralinos are more degenerate, leading to softer decay products, and their branching ratios to  $b$  quarks are enhanced by their Higgsino component. Such issues may have a large impact on chargino and neutralino searches at the Tevatron. This is an important question, but currently the Tevatron reach in the focus point region is unknown.

While the Higgs boson is not a supersymmetric particle, supersymmetry (in its economical implementations) restricts its mass, and so Higgs boson searches also have an important impact on supersymmetric models. For LEP II, the ultimate exclusion limit, barring a discovery, is expected to be  $m_h > 115 \text{ GeV}$ . At the Tevatron, the  $3\sigma$  ( $5\sigma$ ) Higgs boson discovery reach for  $10 \text{ fb}^{-1}$  is  $m_h \lesssim 100$  (120) GeV [128–130]. The Higgs boson mass, unlike all other

TABLE IV. Constraints on supersymmetric models used in Figs. 18 and 19. We also list experiments likely to reach these sensitivities before 2006.

Observable	Type	Bound	Experiment(s)
$\tilde{\chi}^+ \tilde{\chi}^-$	Collider	$m_{\tilde{\chi}}^\pm > 100 \text{ GeV}$	LEP: ALEPH, DELPHI, L3, OPAL
$\tilde{\chi}^\pm \tilde{\chi}^0$	Collider	See Refs. [121,124,126]	Tevatron: CDF, D0
$B \rightarrow X_s \gamma$	Low energy	$ \Delta B(B \rightarrow X_s \gamma)  < 1.2 \times 10^{-4}$	BaBar, BELLE
Muon MDM	Low energy	$ a_\mu^{\text{SUSY}}  < 8 \times 10^{-10}$	Brookhaven E821
$\sigma_{\text{proton}}$	Direct DM	Equation (19)	CDMS, CRESST, GENIUS
$\nu$ from Earth	Indirect DM	$\Phi_\mu^\oplus < 100 \text{ km}^{-2} \text{ yr}^{-1}$	AMANDA, NESTOR, ANTARES
$\nu$ from Sun	Indirect DM	$\Phi_\mu^\odot < 100 \text{ km}^{-2} \text{ yr}^{-1}$	AMANDA, NESTOR, ANTARES
$\gamma$ (gal. center)	Indirect DM	$\Phi_\gamma(1) < 1.5 \times 10^{-10} \text{ cm}^{-2} \text{ s}^{-1}$	GLAST
$\gamma$ (gal. center)	Indirect DM	$\Phi_\gamma(50) < 3 \times 10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$	HESS, CANGAROO III
$e^+$ cosmic rays	Indirect DM	$(S/B)_{\text{max}} < 0.01$	AMS-02

quantities investigated here, is sensitive to the  $A_0$  parameter. As the Higgs boson mass limit rises, models with non-zero  $A_0$ , large  $\tan \beta$ , and  $m_0 \gtrsim 1 \text{ TeV}$  are increasingly favored [22]. However, for natural values of  $A_0$  [22],  $100 \text{ GeV} < m_h < 120 \text{ GeV}$  and so the Higgs boson will be discovered at the Tevatron at  $3\sigma$ , but never at  $5\sigma$ .

Finally, there are many opportunities for discovering supersymmetry in low energy experiments. These include effects in hadronic and leptonic flavor violation, CP violation, proton decay, and electric and magnetic dipole moments. These are discussed more completely in Ref. [24]. Here we will focus on two particularly robust probes:  $B \rightarrow X_s \gamma$  and the anomalous magnetic moment of the muon.

The best current measurements of  $B \rightarrow X_s \gamma$  from CLEO and ALEPH can be combined in a weighted average of  $B(B \rightarrow X_s \gamma)_{\text{exp}} = (3.14 \pm 0.48) \times 10^{-4}$  [131]. These measurements will be improved at the  $B$  factories, where large samples of  $B$  mesons will greatly reduce statistical errors. However, the uncertainty in the theoretical prediction of the standard model,  $B(B \rightarrow X_s \gamma)_{\text{SM}} = (3.29 \pm 0.30) \times 10^{-4}$ , is likely to remain unchanged. By 2006, a conservative estimate is that both theoretical and experimental uncertainties will be  $\sim 0.3 \times 10^{-4}$ . Combining them linearly, the  $2\sigma$  limit will be  $2.1 \times 10^{-4} < B(B \rightarrow X_s \gamma) < 4.5 \times 10^{-4}$ .

The supersymmetric contribution to the muon magnetic dipole moment (MDM)  $a_\mu = \frac{1}{2}(g - 2)_\mu$  is also a robust probe, since it involves only a few (flavor- and CP-conserving) parameters [132]. The world average is  $a_\mu^{\text{exp}} = (116\,592\,05 \pm 45) \times 10^{-10}$  [133] and is consistent with the standard model. However, once data currently being taken is analyzed, the Brookhaven experiment E821 is expected to reduce the uncertainty to  $\Delta a_\mu \sim 4 \times 10^{-10}$  [134]. At present, uncertainties in the standard model prediction are substantial. Assuming these can be reduced, however, a reasonable estimate for future  $2\sigma$  sensitivity is  $a_\mu^{\text{SUSY}} = 8 \times 10^{-10}$ .

In Table IV we present our estimates for sensitivities that will be achieved before the LHC begins operation. The experiments likely to achieve these projections are also listed. Using these estimates, the reach in minimal supergravity parameter space for each mode is given in Figs. 18 and 19. In reading these figures, recall that we have assumed constant local densities in our assessment of dark matter search reaches. If one assumes that the local density is modulated by the thermal relic density, the dark matter reaches outside






FIG. 18. Estimated reaches of various high-energy collider and low-energy precision searches (black), direct dark matter searches (red), and indirect dark matter searches (blue) before the LHC begins operation, for  $\tan \beta = 10$ . The projected sensitivities used are given in Table IV. (The LEP chargino mass bound will marginally extend the bottom and right excluded regions and is omitted.) The shaded regions are as in Fig. 1. The regions probed extend the curves toward the forbidden, green regions. The dark matter reaches are *not* modulated by the thermal relic density. Bounds from photons from the galactic center are highly halo model-dependent; we assume a moderate halo profile parameter  $\bar{J} = 500$ . (See text.)




FIG. 19. As in Fig. 18, but for  $\tan \beta = 50$ . Here the  $\Phi_\gamma^1$  probe is sensitive to all of the parameter space shown and so its limit contour does not appear.

the shaded regions should be suitably diminished. Within the shaded regions, however, our analysis applies without modification.

Several striking features emerge from Figs. 18 and 19. First, we see that, within the minimal supergravity framework, nearly all of the cosmologically preferred models will be probed by at least one experiment. This is strictly true for  $\tan\beta = 10$ . For  $\tan\beta = 50$ , some of the preferred region escapes all probes, but this requires  $M_{1/2} \gtrsim 450$  GeV and  $m_0 \gtrsim 1.5$  TeV, and requires significant fine-tuning of the electroweak scale. In the most natural regions, all models in which neutralinos form a significant fraction of dark matter will yield some signal before the LHC begins operation.

Also noteworthy is the complementarity of traditional particle physics searches and indirect dark matter searches. Collider searches require, of course, light superpartners. High precision probes at low energy also require light superpartners, as the virtual effects of superpartners quickly decouple as they become heavy. Thus, the LEP and Tevatron reaches are confined to the lower left-hand corner, as are, to a lesser extent, the searches for deviations in  $B \rightarrow X_s \gamma$  and  $a_\mu$ . These bounds, and all others of this type, are easily satisfied in the focus point models with large  $m_0$ , and indeed this is one of the virtues of these models. However, in the focus point models, *all* of the indirect searches are maximally sensitive, as the dark matter contains a significant Higgsino component. Direct dark matter probes share features with both traditional and indirect searches, and have sensitivity in both regions. It is only by combining all of these experiments, that the preferred region may be completely explored.<sup>3</sup>

Finally, these results have implications for future colliders. In the cosmologically preferred regions of parameter space with  $0.1 < \Omega_\chi h^2 < 0.3$ , all models with charginos or sleptons lighter than 300 GeV will produce observable signals in at least one experiment. This is evident for  $\tan\beta = 10$  and 50 in Figs. 18 and 19. In Fig. 20, we vary  $\tan\beta$ , fixing  $M_{1/2}$  to 400 GeV, which roughly corresponds to 300 GeV charginos. We see that the preferred region is probed for any choice of  $\tan\beta$ . (For extremely low  $\tan\beta$  and  $m_0$ , there appears to be a region that is not probed. However, this is excluded by current Higgs mass limits for  $A_0 = 0$ . These limits might be evaded if  $A_0$  is also tuned to some extreme value, but in this case, top squark searches in Run II of the Tevatron will provide an additional constraint.)

These results imply that if any superpartners are to be within reach of a 500 GeV lepton collider, some hint of supersymmetry must be seen before the LHC begins collecting data. This conclusion is independent of naturalness considerations. While our quantitative analysis is confined to minimal supergravity, we expect this result to be valid more generally. For moderate values of  $\tan\beta$ , if the dark matter is made up of neutralinos, they must either be light, Bino-like, or a gaugino-Higgsino mixture. If they are light, charginos will be discovered. If they are Bino-like, light sfermions are required to mediate their annihilation, and there will be anomalies in low energy precision measurements. And if they are a gaugino-Higgsino

---

<sup>3</sup>Note that the complementarity referred to here is not the commonly recognized one, which concerns the mass of the neutralino. It is well-known that some indirect searches are effective even for LSP masses in the TeV range, well beyond the range of colliders. However, such models are highly unnatural, and they have not been considered here.




FIG. 20. As in Fig. 18, but in the  $(m_0, \tan \beta)$  plane for fixed  $M_{1/2} = 400$  GeV,  $A_0 = 0$ , and  $\mu > 0$ . The regions probed are toward the green regions, except for  $\Phi_\gamma^{50}$ , where it is between the two contours. The top excluded region is forbidden by limits on the CP-odd Higgs scalar mass.

mixture, at least one indirect dark matter search will see a signal. For large  $\tan \beta$ , low energy probes become much more effective and again there is sensitivity to all probe superpartner spectra with light superpartners. Thus it appears, on qualitative grounds, that all models in which the scalar masses are not widely separated, and the charginos are not extravagantly heavy, will be accessible prior to LHC operation.

## VII. CONCLUSIONS

In this paper, we have examined a wide variety of indirect dark matter detection signals. These include neutrinos from annihilation of dark matter in the cores of the Earth and Sun, continuum gamma rays from annihilation in the galactic center, and positron excesses in cosmic rays from annihilation in the local solar neighborhood. In each case, the experimental landscape will be transformed in the next few years by experiments that are running or being mounted. We have tabulated the salient features and reaches of some of the most promising experiments in the previous sections.

We have evaluated the prospects for dark matter detection in the framework of minimal supergravity. This framework incorporates many of the most compelling features of supersymmetry. Previously, this framework has been thought to predict a Bino-like LSP. That severely limited its utility for dark matter studies. However, recent work has made it clear that gaugino-Higgsino mixtures and even Higgsino-like LSPs are also quite naturally realized in minimal supergravity. We have been careful to include the full range of possibilities, with important (and positive) implications for future dark matter searches.

Let us note in passing that in our parametrization of experimental probes, the case of no-scale supergravity [135], recently revived in the context of gaugino-mediated supersymmetry

breaking [136–139], can be regarded as the special case  $m_0 = 0$ . Experimental probes of these models are simply evaluated by restricting to the  $m_0 = 0$  axis. Several experiments, notably the trilepton Tevatron search, direct dark matter searches, and Brookhaven experiment E821 will have the power to confirm or exclude this possibility in the near future.

We have concentrated here on discovery signals. If a signal is confirmed, precision measurements may allow experiments to determine dark matter properties. For example, as has been noted in the literature, the neutralino’s mass may be determined by the angular spread of the signal in neutrino telescopes. The energy spectrum of gamma rays or positrons signals may provide similar information. The gaugino-ness may also be constrained; indeed, the existence of a significant signal in itself would constitute evidence in favor of mixed gaugino-Higgsinos.

The simplicity of minimal supergravity allows us to compare the reaches of a great variety of probes. We summarize by collecting several of our main conclusions:

- Bino-like dark matter leads to suppressed rates for all indirect dark matter signals. In this case, unless the neutralino is extremely light (near current bounds), all indirect signals are beyond detection for the foreseeable future.
- Higgsino-like dark matter cannot yield cosmologically interesting relic densities in a straightforward way. Studies that assume Higgsino-like dark matter exaggerate the power of indirect searches.
- Mixed gaugino-Higgsino dark matter gives both relic densities in the preferred range  $0.1 \lesssim \Omega_\chi h^2 \lesssim 0.3$  and detectable signals. Such dark matter is naturally present in focus point models, which are favored by low-energy constraints.
- Experiments that are running or underway will transform the prospects for indirect dark matter detection. Among the most promising experiments are the neutrino telescopes AMANDA, NESTOR, and ANTARES; the gamma ray telescopes MAGIC, HESS, and CANGAROO III, and the satellite detector GLAST; and AMS-02, the anti-particle/anti-matter search aboard the International Space Station. For mixed gaugino-Higgsino dark matter, these experiments will be sensitive to nearly all models with cosmologically interesting neutralino relic densities, and are competitive with next-generation direct search experiments, such as CDMS (Soudan) and CRESST.
- The various indirect searches rely on different sources of neutralino annihilation (cores of the Earth or Sun, galactic center, local solar neighborhood) and so are sensitive to different assumptions. In addition, some signals, particularly the continuum photons, will be difficult to identify unambiguously as a dark matter signal. Without actual data and detailed analyses, it is difficult to make a more precise statement. However, we have seen that many experiments are sensitive to the same supersymmetric models, and given the underlying uncertainties, redundancy is clearly a virtue.
- Indirect searches are complementary to traditional particle searches. Separately they probe only portions of the cosmologically preferred model space. Combined, essentially all cosmologically preferred models will produce at least a hint of a signal in one of these experiments *before* the LHC begins operation.

- In minimal supergravity models with  $0.1 < \Omega_\chi h^2 < 0.3$ , *if there is no hint of supersymmetry before the LHC begins operation, no superpartners will be within reach of a 500 GeV lepton collider*. Our arguments are independent of naturalness considerations, and their qualitative structure suggests that similar conclusions will remain valid in alternative frameworks.

## ACKNOWLEDGMENTS

We thank P. Blasi, L. Hui, Z. Ligeti and U. Nierste for helpful discussions and readings of the manuscript, and I. Moskalenko for correspondence. This work was supported in part by the Department of Energy under contracts DE-FG02-90ER40542 and DE-AC02-76CH03000, and by the National Science Foundation under grant PHY-9513835. J.L.F. acknowledges the support of a Frank and Peggy Taplin Fellowship.

## REFERENCES

- [1] For a recent review, see J. R. Primack, “Cosmological Parameters 2000,” astro-ph/0007187.
- [2] K. Freese, B. D. Fields and D. S. Graff, “Death of Stellar Baryonic Dark Matter,” in *Proceedings of the MPA/ESO Workshop on the First Stars*, Garching, Germany, 4-6 August 1999, astro-ph/0002058.
- [3] H. Goldberg, “Constraint on the photino mass from cosmology,” Phys. Rev. Lett. **50**, 1419 (1983).
- [4] J. Ellis, J. S. Hagelin, D. V. Nanopoulos and M. Srednicki, “Search For Supersymmetry At The  $\bar{p}p$  Collider,” Phys. Lett. **B127**, 233 (1983).
- [5] LEPSUSYWG, ALEPH, DELPHI, L3 and OPAL experiments, Note LEPSUSYWG/99-03.1.
- [6] J. L. Feng, K. T. Matchev and F. Wilczek, “Neutralino dark matter in focus point supersymmetry,” Phys. Lett. **B482**, 388 (2000) [hep-ph/0004043].
- [7] For excellent reviews, see, *e.g.*, M. Drees, “Particle dark matter physics: An update,” Pramana **51**, 87 (1998) [hep-ph/9804231]; F. Halzen, “Lectures on neutrino astronomy: Theory and experiment,” astro-ph/9810368; J. Ellis, “Particles and cosmology: Learning from cosmic rays,” astro-ph/9911440. L. Bergström, “Non-baryonic dark matter: Observational evidence and detection methods,” hep-ph/0002126.
- [8] A. E. Faraggi, K. A. Olive and M. Pospelov, “Probing the desert with ultra-energetic neutrinos from the sun,” Astropart. Phys. **13**, 31 (2000) [hep-ph/9906345].
- [9] A. Corsetti and P. Nath, “Gaugino mass nonuniversality and dark matter in SUGRA, strings and D brane models,” hep-ph/0003186.
- [10] G. F. Giudice and E. Roulet, “Energetic Neutrinos From Supersymmetric Dark Matter,” Nucl. Phys. **B316**, 429 (1989).
- [11] G. B. Gelmini, P. Gondolo and E. Roulet, “Neutralino dark matter searches,” Nucl. Phys. **B351**, 623 (1991).
- [12] R. Gandhi, J. L. Lopez, D. V. Nanopoulos, K. Yuan and A. Zichichi, “Scrutinizing supergravity models through neutrino telescopes,” Phys. Rev. **D49**, 3691 (1994) [astro-ph/9309048].
- [13] E. Diehl, G. L. Kane, C. Kolda and J. D. Wells, “Theory, phenomenology, and prospects for detection of supersymmetric dark matter,” Phys. Rev. **D52**, 4223 (1995) [hep-ph/9502399].
- [14] A. Corsetti and P. Nath, “Out-going muon flux from neutralino annihilation in the sun and the earth in supergravity unification,” Int. J. Mod. Phys. **A15**, 905 (2000) [hep-ph/9904497].
- [15] S. Dimopoulos, S. Raby and F. Wilczek, “Supersymmetry And The Scale Of Unification,” Phys. Rev. **D24**, 1681 (1981).
- [16] L. Bergström and P. Gondolo, “Limits on direct detection of neutralino dark matter from  $b \rightarrow s\gamma$  decays,” Astropart. Phys. **5**, 263 (1996) [hep-ph/9510252].
- [17] L. Bergström, J. Edsjö and P. Gondolo, “Indirect neutralino detection rates in neutrino telescopes,” Phys. Rev. **D55**, 1765 (1997) [hep-ph/9607237].
- [18] L. Bergström, J. Edsjö and P. Gondolo, “Indirect detection of dark matter in km-size neutrino telescopes,” Phys. Rev. **D58**, 103519 (1998) [hep-ph/9806293].
- [19] L. Bergström, P. Ullio and J. H. Buckley, “Observability of gamma rays from dark matter neutralino annihilations in the Milky Way halo,” Astropart. Phys. **9**, 137 (1998) [astro-ph/9712318].
- [20] E. A. Baltz, C. Briot, P. Salati, R. Taillet and J. Silk, “Detection of neutralino annihilation photons from external galaxies,” Phys. Rev. **D61**, 023514 (2000) [astro-ph/9909112].

- [21] J. L. Feng, K. T. Matchev and T. Moroi, “Multi-TeV scalars are natural in minimal supergravity,” *Phys. Rev. Lett.* **84**, 2322 (2000) [hep-ph/9908309].
- [22] J. L. Feng, K. T. Matchev and T. Moroi, “Focus points and naturalness in supersymmetry,” *Phys. Rev.* **D61**, 075005 (2000) [hep-ph/9909334].
- [23] J. L. Feng, K. T. Matchev and T. Moroi, “Naturalness re-examined: Implications for supersymmetry searches,” in *Proceedings of the 7th International Symposium on Particles, Strings and Cosmology (PASCOS 99)*, Granlibakken, Tahoe City, California, 10-16 Dec 1999, hep-ph/0003138.
- [24] K. Matchev, talk given at PHENO'00, Madison, WI, 18 April 2000; J. Feng, talk given at the Theoretical Institute on SUSY and Higgs 2000, Argonne, 12 May 2000; J. L. Feng and K. T. Matchev, in preparation.
- [25] D. M. Pierce, J. A. Bagger, K. Matchev and R. Zhang, “Precision corrections in the minimal supersymmetric standard model,” *Nucl. Phys.* **B491**, 3 (1997) [hep-ph/9606211].
- [26] S. P. Martin and M. T. Vaughn, “Two loop renormalization group equations for soft supersymmetry breaking couplings,” *Phys. Rev.* **D50**, 2282 (1994) [hep-ph/9311340]; I. Jack and D. R. Jones, “Soft supersymmetry breaking and finiteness,” *Phys. Lett.* **B333**, 372 (1994) [hep-ph/9405233]; Y. Yamada, “Two loop renormalization group equations for soft SUSY breaking scalar interactions: Supergraph method,” *Phys. Rev.* **D50**, 3537 (1994) [hep-ph/9401241]; I. Jack, D. R. Jones, S. P. Martin, M. T. Vaughn and Y. Yamada, “Decoupling of the epsilon scalar mass in softly broken supersymmetry,” *Phys. Rev.* **D50**, 5481 (1994) [hep-ph/9407291].
- [27] J. Bagger, K. Matchev and D. Pierce, “Precision corrections to supersymmetric unification,” *Phys. Lett.* **B348**, 443 (1995) [hep-ph/9501277].
- [28] S. Mizuta, D. Ng and M. Yamaguchi, “Phenomenological aspects of supersymmetric standard models without grand unification,” *Phys. Lett.* **B300**, 96 (1993) [hep-ph/9210241].
- [29] T. Moroi and L. Randall, “Wino cold dark matter from anomaly-mediated SUSY breaking,” *Nucl. Phys.* **B570**, 455 (2000) [hep-ph/9906527].
- [30] R. Barbieri and G. F. Giudice, “Upper Bounds On Supersymmetric Particle Masses,” *Nucl. Phys.* **B306**, 63 (1988).
- [31] M. Carena, M. Olechowski, S. Pokorski and C. E. Wagner, “Electroweak symmetry breaking and bottom - top Yukawa unification,” *Nucl. Phys.* **B426**, 269 (1994) [hep-ph/9402253].
- [32] S. Mizuta and M. Yamaguchi, “Coannihilation effects and relic abundance of Higgsino dominant  $LSP_s$ ,” *Phys. Lett.* **B298**, 120 (1993) [hep-ph/9208251].
- [33] J. Ellis, T. Falk and K. A. Olive, “Neutralino stau coannihilation and the cosmological upper limit on the mass of the lightest supersymmetric particle,” *Phys. Lett.* **B444**, 367 (1998) [hep-ph/9810360].
- [34] J. Ellis, T. Falk, K. A. Olive and M. Srednicki, “Calculations of neutralino stau coannihilation channels and the cosmologically relevant region of MSSM parameter space,” *Astropart. Phys.* **13**, 181 (2000) [hep-ph/9905481].
- [35] M. E. Gomez, G. Lazarides and C. Pallis, “Yukawa unification,  $b \rightarrow s\gamma$  and bino-stau coannihilation,” hep-ph/0004028.
- [36] P. H. Chankowski, J. Ellis, M. Olechowski and S. Pokorski, “Haggling over the fine-tuning price of LEP,” *Nucl. Phys.* **B544**, 39 (1999) [hep-ph/9808275].
- [37] K. Freese, “Can Scalar Neutrinos Or Massive Dirac Neutrinos Be The Missing Mass?,” *Phys. Lett.* **B167**, 295 (1986).

- [38] L. M. Krauss, M. Srednicki and F. Wilczek, “Solar System Constraints And Signatures For Dark Matter Candidates,” *Phys. Rev.* **D33**, 2079 (1986).
- [39] T. K. Gaisser, G. Steigman and S. Tilav, “Limits On Cold Dark Matter Candidates From Deep Underground Detectors,” *Phys. Rev.* **D34**, 2206 (1986).
- [40] A. Gould, J. A. Frieman and K. Freese, “Probing The Earth With Wimps,” *Phys. Rev.* **D39**, 1029 (1989).
- [41] W. H. Press and D. N. Spergel, “Capture by the sun of a galactic population of weakly interacting, massive particles,” *Astrophys. J.* **296**, 679 (1985).
- [42] J. Silk, K. Olive and M. Srednicki, “The photino, the sun, and high-energy neutrinos,” *Phys. Rev. Lett.* **55**, 257 (1985).
- [43] M. Srednicki, K. A. Olive and J. Silk, “High-Energy Neutrinos From The Sun And Cold Dark Matter,” *Nucl. Phys.* **B279**, 804 (1987).
- [44] J. S. Hagelin, K. W. Ng and K. A. Olive, “A High-Energy Neutrino Signature From Supersymmetric Relics,” *Phys. Lett.* **B180**, 375 (1986).
- [45] K. Ng, K. A. Olive and M. Srednicki, “Dark Matter Induced Neutrinos From The Sun: Theory Versus Experiment,” *Phys. Lett.* **B188**, 138 (1987).
- [46] J. Ellis and R. A. Flores, “Realistic Predictions For The Detection Of Supersymmetric Dark Matter,” *Nucl. Phys.* **B307**, 883 (1988).
- [47] G. Jungman, M. Kamionkowski and K. Griest, “Supersymmetric dark matter,” *Phys. Rept.* **267**, 195 (1996) [hep-ph/9506380];  
<http://t8web.lanl.gov/people/jungman/neut-package.html>.
- [48] L. Bergström, T. Damour, J. Edsjö, L. M. Krauss and P. Ullio, “Implications of a new solar system population of neutralinos on indirect detection rates,” *JHEP* **9908**, 010 (1999) [hep-ph/9905446].
- [49] A. Gould and S. M. Khairul Alam, “Can heavy WIMPs be captured by the earth?,” *astro-ph/9911288*.
- [50] A. Gould, “Wimp Distribution In And Evaporation From The Sun,” *Astrophys. J.* **321**, 560 (1987).
- [51] A. Gould, “Resonant Enhancements In Wimp Capture By The Earth,” *Astrophys. J.* **321**, 571 (1987).
- [52] A. Gould, “Direct And Indirect Capture Of Wimps By The Earth,” *Astrophys. J.* **328**, 919 (1988).
- [53] S. Ritz and D. Seckel, “Detailed Neutrino Spectra From Cold Dark Matter Annihilations In The Sun,” *Nucl. Phys.* **B304**, 877 (1988).
- [54] G. Jungman and M. Kamionkowski, “Neutrinos from particle decay in the sun and earth,” *Phys. Rev.* **D51**, 328 (1995) [hep-ph/9407351].
- [55] M. Kamionkowski, “Energetic neutrinos from heavy neutralino annihilation in the sun,” *Phys. Rev.* **D44**, 3021 (1991).
- [56] A. Bottino, V. de Alfaro, N. Fornengo, G. Mignola and M. Pignone, “Indirect search for neutralinos at neutrino telescopes,” *Phys. Lett.* **B265**, 57 (1991).
- [57] A. Bottino, N. Fornengo, G. Mignola and L. Moscoso, “Signals of neutralino dark matter from earth and sun,” *Astropart. Phys.* **3**, 65 (1995) [hep-ph/9408391].
- [58] V. Berezhinsky, A. Bottino, J. Ellis, N. Fornengo, G. Mignola and S. Scopel, “Searching for relic neutralinos using neutrino telescopes,” *Astropart. Phys.* **5**, 333 (1996) [hep-ph/9603342].
- [59] K. Freese and M. Kamionkowski, “Indirect detection of a light Higgsino motivated by collider



- data,” Phys. Rev. **D55**, 1771 (1997) [hep-ph/9609370].
- [60] A. Bottino, N. Fornengo, G. Mignola, M. Olechowski and S. Scopel, “Perspectives for detection of a higgsino-like relic neutralino,” Astropart. Phys. **6**, 395 (1997) [astro-ph/9611030].
  - [61] A. Bottino, F. Donato, N. Fornengo and S. Scopel, “Combining the data of annual modulation effect in WIMP direct detection with measurements of WIMP indirect searches,” Astropart. Phys. **10**, 203 (1999) [hep-ph/9809239].
  - [62] A. Bottino, F. Donato, N. Fornengo and S. Scopel, “Further investigation of a relic neutralino as a possible origin of an annual-modulation effect in WIMP direct search,” hep-ph/0001309.
  - [63] *Proceedings of the 26th International Cosmic Ray Conference (ICRC 99)*, Salt Lake City, Utah, 17-25 August 1999, <http://krusty.physics.utah.edu/icrc1999/proceedings.html>.
  - [64] Baksan Collaboration, S. P. Mikheyev *et al.*, talk given at the 8th International Workshop on Neutrino Telescopes, Venice, Italy, 23-26 February 1999; O. V. Suvorova, talk given at the 2nd International Conference on Physics Beyond the Standard Model: Beyond the Desert 99, Ringberg Castle, Tegernsee, Germany, 6-12 June 1999, hep-ph/9911415.
  - [65] Kamiokande Collaboration, M. Mori *et al.*, Phys. Rev. **D48**, 5505 (1993).
  - [66] MACRO Collaboration, M. Ambrosio *et al.*, Phys. Rev. **D60**, 082002 (1999), hep-ex/9812020; MACRO Collaboration, T. Montaruli *et al.*, HE.5.1.03 in Ref. [63], hep-ex/9905021.
  - [67] Super-Kamiokande Collaboration, A. Okada *et al.*, “Searches for astronomical neutrino sources and WIMPs with Super-Kamiokande,” astro-ph/0007003.
  - [68] Baikal Collaboration, V. A. Balkanov *et al.*, in *The proceedings of the 6th International Workshop on Topics in Astroparticle and Underground Physics (TAUP 99)*, Paris, France, 6-10 September 1999, astro-ph/0001151.
  - [69] AMANDA Collaboration, E. Dalberg *et al.*, HE.5.3.06 in Ref. [63]; AMANDA Collaboration, A. Karle *et al.*, HE.4.2.05 in Ref. [63].
  - [70] AMANDA Collaboration, C. Spiering *et al.*, talk given at the 8th International Workshop on Neutrino Telescopes, Venice, Italy, 23-26 February 1999, astro-ph/9906205.
  - [71] NESTOR Collaboration, L. K. Resvanis *et al.*, talk given at the 8th International Workshop on Neutrino Telescopes, Venice, Italy, 23-26 February 1999.
  - [72] ANTARES Collaboration, J. R. Hubbard *et al.*, HE.6.3.03 in Ref. [63]; ANTARES Collaboration, L. Moscoso *et al.*, HE.6.3.04 in Ref. [63].
  - [73] L. Bergström, J. Edsjö and M. Kamionkowski, “Astrophysical-neutrino detection with angular and energy resolution,” Astropart. Phys. **7**, 147 (1997) [astro-ph/9702037].
  - [74] M. Urban, A. Bouquet, B. Degrange, P. Fleury, J. Kaplan, A. L. Melchior and E. Pare, “Searching for TeV dark matter by atmospheric Cerenkov techniques,” Phys. Lett. **B293**, 149 (1992) [hep-ph/9208255].
  - [75] V. S. Berezinsky, A. V. Gurevich and K. P. Zybin, “Distribution of dark matter in the galaxy and the lower limits for the masses of supersymmetric particles,” Phys. Lett. **B294**, 221 (1992).
  - [76] V. Berezinsky, A. Bottino and G. Mignola, “High-energy gamma radiation from the galactic center due to neutralino annihilation,” Phys. Lett. **B325**, 136 (1994) [hep-ph/9402215].
  - [77] L. Bergström, J. Edsjö and P. Ullio, “Possible Indications of a Clumpy Dark Matter Halo,” Phys. Rev. **D58**, 083507 (1998) [astro-ph/9804050].
  - [78] L. Bergström, J. Edsjö, P. Gondolo and P. Ullio, “Clumpy neutralino dark matter,” Phys. Rev. **D59**, 043506 (1999) [astro-ph/9806072].
  - [79] P. Gondolo and J. Silk, “Dark matter annihilation at the galactic center,” Phys. Rev. Lett.

- 83**, 1719 (1999) [astro-ph/9906391].
- [80] L. Bergström and P. Ullio, “Full one-loop calculation of neutralino annihilation into two photons,” Nucl. Phys. **B504**, 27 (1997) [hep-ph/9706232].
  - [81] Z. Bern, P. Gondolo and M. Perelstein, “Neutralino annihilation into two photons,” Phys. Lett. **B411**, 86 (1997) [hep-ph/9706538].
  - [82] P. Ullio and L. Bergström, “Neutralino annihilation into a photon and a Z boson,” Phys. Rev. **D57**, 1962 (1998) [hep-ph/9707333].
  - [83] V. S. Berezinsky, A. Bottino and V. de Alfaro, “Is it possible to detect the gamma ray line from neutralino-neutralino annihilation?,” Phys. Lett. **B274**, 122 (1992).
  - [84] J. F. Navarro, C. S. Frenk and S. D. White, “The Structure of Cold Dark Matter Halos,” Astrophys. J. **462**, 563 (1996) [astro-ph/9508025].
  - [85] P. Gondolo, “Either neutralino dark matter or cuspy dark halos,” hep-ph/0002226.
  - [86] H. U. Bengtsson, P. Salati and J. Silk, “Quark Flavors And The Gamma Ray Spectrum From Halo Dark Matter Annihilations,” Nucl. Phys. **B346**, 129 (1990).
  - [87] <http://coss.gsfc.nasa.gov/egret>.
  - [88] STACEE Collaboration, M. C. Chantell *et al.*, Nucl. Instrum. Meth. **A408**, 468 (1998), astro-ph/9704037; R. Ong, talk at the 6th GeV-TeV Gamma Ray Astrophysics Workshop, Snowbird, Utah, 13-16 August 1999, <http://hep.uchicago.edu/~stacee>.
  - [89] CELESTE and CAT Collaborations, M. de Naurois *et al.*, OG.4.3.06 in Ref. [63]; <http://www.cenbg.in2p3.fr/Astroparticule/celeste/e-index.html>.
  - [90] ARGO-YBJ Collaboration, M. Iacovacci *et al.*, Nucl. Phys. Proc. Suppl. **85**, 338 (2000), hep-ex/9908060; <http://www1.na.infn.it/wsubnucl/cosm/argo/argo.html>.
  - [91] MAGIC Collaboration, M. Martinez *et al.*, OG.4.3.08 in Ref. [63]; <http://hegral.mppmu.mpg.de>.
  - [92] AGILE Collaboration, A. Morselli *et al.*, OG.4.2.06 in Ref. [63]; <http://www.ifctr.mi.cnr.it/Agile>.
  - [93] HESS Collaboration, A. Kohnle *et al.*, OG.4.3.13 and OG.4.3.24 in Ref. [63]; <http://www-hfm.mpi-hd.mpg.de/HESS>.
  - [94] R. Battiston, M. Biasini, E. Fiandrini, J. Petrakis, and M. H. Salamon, Astropart. Phys. **13**, 51 (2000), astro-ph/9909432; R. Battiston, talk given at the 6th GeV-TeV Gamma Ray Astrophysics Workshop, Snowbird, Utah, 13-16 August 1999, astro-ph/9911241.
  - [95] CANGAROO Collaboration, M. Mori *et al.*, talk given at the 6th GeV-TeV Gamma Ray Astrophysics Workshop, Snowbird, Utah, 13-16 August 1999; <http://icrhp9.icrr.u-tokyo.ac.jp>.
  - [96] VERITAS Collaboration, S. M. Bradbury *et al.*, OG.4.3.28 in Ref. [63], astro-ph/9907248; <http://earth.physics.purdue.edu/veritas>.
  - [97] GLAST Collaboration, A. Moiseev *et al.*, HE.5.1.02 in Ref. [63]; <http://www-glast.stanford.edu>.
  - [98] P. Chardonnet, P. Salati, J. Silk, I. Grenier and G. Smoot, “The Gamma ray diffuse background and Cerenkov telescopes,” Ap. J. **454**, 774 (1995).
  - [99] EGRET Collaboration, S. D. Hunter *et al.*, “EGRET Observations of the Diffuse Gamma-Ray Emission from the Galactic Plane,” Ap. J. **481**, 205 (1997).
  - [100] A. J. Tylka, “Cosmic Ray Positrons From Annihilation Of Weakly Interacting Massive Particles In The Galaxy,” Phys. Rev. Lett. **63**, 840 (1989).
  - [101] M. S. Turner and F. Wilczek, “Positron Line Radiation From Halo Wimp Annihilations As A Dark Matter Signature,” Phys. Rev. **D42**, 1001 (1990).

- [102] M. Kamionkowski and M. S. Turner, “A Distinctive positron feature from heavy WIMP annihilations in the galactic halo,” *Phys. Rev.* **D43**, 1774 (1991).
- [103] E. A. Baltz and J. Edsjo, “Positron propagation and fluxes from neutralino annihilation in the halo,” *Phys. Rev.* **D59**, 023511 (1999) [astro-ph/9808243].
- [104] I. V. Moskalenko and A. W. Strong, “Positrons from particle dark-matter annihilation in the galactic halo: Propagation Green’s functions,” *Phys. Rev.* **D60**, 063003 (1999) [astro-ph/9905283].
- [105] F. W. Stecker and A. J. Tylka, “The Cosmic Ray Anti-Proton Spectrum From Dark Matter Annihilation And Its Astrophysical Implications: A New Look,” *Ap. J.* **336**, L51 (1989).
- [106] P. Chardonnet, G. Mignola, P. Salati and R. Taillet, “Galactic diffusion and the antiproton signal of supersymmetric dark matter,” *Phys. Lett.* **B384**, 161 (1996) [astro-ph/9606174].
- [107] L. Bergström, J. Edsjö and P. Ullio, “Cosmic antiprotons as a probe for supersymmetric dark matter?,” astro-ph/9902012.
- [108] J. W. Bieber, R. A. Burger, R. Engel, T. K. Gaisser, S. Roesler and T. Stanev, “Antiprotons at solar maximum,” *Phys. Rev. Lett.* **83**, 674 (1999) [astro-ph/9903163].
- [109] F. Donato, N. Fornengo and P. Salati, “Antideuterons as a signature of supersymmetric dark matter,” hep-ph/9904481.
- [110] R. J. Protheroe, “On The Nature Of The Cosmic Ray Positron Spectrum,” *Astrophys. J.* **254**, 391 (1982).
- [111] HEAT Collaboration, S. W. Barwick *et al.*, “Cosmic ray positrons at high-energies: A New measurement,” *Phys. Rev. Lett.* **75**, 390 (1995) [astro-ph/9505141]; M. A. DuVernois *et al.*, OG.1.1.14 in Ref. [63].
- [112] WIZARD-CAPRICE Collaboration, M. Boezio *et al.*, “The cosmic-ray electron and positron spectra measured at 1 AU during solar minimum activity,” *Astrophys. J.* **532**, 653 (2000); M. Boezio *et al.*, OG.1.1.16 in Ref. [63]; M. Boezio, Ph. D. thesis, Royal Institute of Technology, Stockholm, Sweden (1998).
- [113] PAMELA Collaboration, OG.4.2.04 in Ref. [63]; [http://wizard.roma2.infn.it/pamela/fram\\_des.htm](http://wizard.roma2.infn.it/pamela/fram_des.htm).
- [114] AMS Collaboration, S. Ahlen *et al.*, *Nucl. Instrum. Meth.* **A350**, 351 (1994); B. Alpat *et al.*, “The AMS (Alpha Magnetic Spectrometer) Experiment on the International Space Station,” talk given at the 8th Pisa meeting on Advanced Detectors: Frontier Detectors for Frontier Physics, 21-27 May 2000; <http://ams.cern.ch/AMS>.
- [115] DAMA Collaboration, R. Bernabei *et al.*, “Search for WIMP annual modulation signature: Results from DAMA / NaI-3 and DAMA / NaI-4 and the global combined analysis,” *Phys. Lett.* **B480**, 23 (2000).
- [116] L. Baudis and H. V. Klapdor-Kleingrothaus, “Direct detection of nonbaryonic dark matter,” astro-ph/0003434.
- [117] CDMS Collaboration, R. W. Schnee *et al.*, “Results and status of the Cryogenic Dark Matter Search (CDMS),” *Phys. Rept.* **307**, 283 (1998).
- [118] CRESST Collaboration, M. Bravin *et al.*, “The CRESST dark matter search,” *Astropart. Phys.* **12**, 107 (1999) [hep-ex/9904005].
- [119] GENUIS Collaboration, H. V. Klapdor-Kleingrothaus *et al.*, “GENIUS: A supersensitive germanium detector system for rare events,” hep-ph/9910205.
- [120] SUGRA Working Group Collaboration, S. Abel *et al.*, “Report of the SUGRA working group for run II of the Tevatron,” hep-ph/0003154.

- [121] K. T. Matchev and D. M. Pierce, “Supersymmetry reach of the Tevatron via trilepton, like-sign dilepton and dilepton plus tau jet signatures,” Phys. Rev. **D60**, 075004 (1999) [hep-ph/9904282].
- [122] H. Baer, M. Drees, F. Paige, P. Quintana and X. Tata, “Trilepton signal for supersymmetry at the Fermilab Tevatron revisited,” Phys. Rev. **D61**, 095007 (2000) [hep-ph/9906233].
- [123] V. Barger and C. Kao, “Trilepton signature of minimal supergravity at the upgraded Tevatron,” Phys. Rev. **D60**, 115015 (1999) [hep-ph/9811489].
- [124] K. T. Matchev and D. M. Pierce, “New backgrounds in trilepton, dilepton and dilepton plus tau jet SUSY signals at the Tevatron,” Phys. Lett. **B467**, 225 (1999) [hep-ph/9907505].
- [125] H. Baer, C. Chen, M. Drees, F. Paige and X. Tata, “Supersymmetry reach of Tevatron upgrades: The large  $\tan\beta$  case,” Phys. Rev. **D58**, 075008 (1998) [hep-ph/9802441].
- [126] J. D. Lykken and K. T. Matchev, “Supersymmetry signatures with tau jets at the Tevatron,” Phys. Rev. **D61**, 015001 (2000) [hep-ph/9903238]; “Tau jet signals for supersymmetry at the Tevatron,” hep-ex/9910033.
- [127] B. C. Allanach, J. P. Hetherington, M. A. Parker and B. R. Webber, “Naturalness reach of the Large Hadron Collider in minimal supergravity,” hep-ph/0005186.
- [128] Report of the Higgs Working Group, conveners M. Carena, J. Conway, H. Haber, and J. Hobbs, <http://fnth37.fnal.gov/higgs.html>.
- [129] H. Baer, B. W. Harris and X. Tata, “The reach of CERN LEP2 and Fermilab Tevatron upgrades for Higgs bosons in supersymmetric models,” Phys. Rev. **D59**, 015003 (1999) [hep-ph/9807262].
- [130] M. Carena, S. Mrenna and C. E. Wagner, “MSSM Higgs boson phenomenology at the Tevatron collider,” Phys. Rev. **D60**, 075010 (1999) [hep-ph/9808312].
- [131] A. L. Kagan and M. Neubert, “QCD anatomy of  $B \rightarrow X_s \gamma$  decays,” Eur. Phys. J. **C7**, 5 (1999) [hep-ph/9805303].
- [132] T. Moroi, “The Muon Anomalous Magnetic Dipole Moment in the Minimal Supersymmetric Standard Model,” Phys. Rev. **D53**, 6565 (1996) [hep-ph/9512396].
- [133] R. Carey, talk given at the XXXth International Conference on High Energy Physics (ICHEP00), Osaka, Japan, 27 July – 2 August 2000.
- [134] E821 Collaboration, M. Grosse Perdekamp *et al.*, “Status of the g-2 experiment at BNL,” Nucl. Phys. Proc. Suppl. **76**, 253 (1999).
- [135] A. B. Lahanas and D. V. Nanopoulos, “The Road To No Scale Supergravity,” Phys. Rept. **145**, 1 (1987).
- [136] D. E. Kaplan, G. D. Kribs and M. Schmaltz, “Supersymmetry breaking through transparent extra dimensions,” Phys. Rev. **D62**, 035010 (2000) [hep-ph/9911293].
- [137] Z. Chacko, M. A. Luty, A. E. Nelson and E. Ponton, “Gaugino mediated supersymmetry breaking,” JHEP **0001**, 003 (2000) [hep-ph/9911323].
- [138] M. Schmaltz and W. Skiba, “Minimal gaugino mediation,” hep-ph/0001172.
- [139] M. Schmaltz and W. Skiba, “The superpartner spectrum of gaugino mediation,” hep-ph/0004210.